



## PROTOCOL

**A randomised, double-blind, placebo-controlled, Phase III trial to determine the efficacy and safety of inhaled SNG001 for the treatment of patients hospitalised due to moderate COVID-19**

**Short Title:** SARS-CoV-2: Phase III TRial of Inhaled  
INTERferon-β Therapy

**Trial Name:** SPRINTER

**Compound:** IFN-β1a (for inhalation)

**Compound Name:** SNG001

**EudraCT Number:** 2020-004743-83

**Protocol Number:** SG018

**Version and Date:** 3, 21 December 2020

### Previous Document History:

Document	Date
Version 2	20 November 2020
Version 1	15 October 2020

**This document is the property of Synairgen Research Ltd and may not be used, divulged, published, or otherwise disclosed without written consent of Synairgen Research Ltd.**

**SYNAIRGEN RESEARCH LTD**

**CLINICAL STUDY PROTOCOL SG018**

**Study Title:** A randomised, double-blind, placebo-controlled, Phase III trial to determine the efficacy and safety of inhaled SNG001 for the treatment of patients hospitalised due to moderate COVID-19

**Protocol Number:** SG018

**Investigational Product:** SNG001 (IFN-β1a for inhalation)

**Sponsor:** Synairgen Research Ltd  
Level F (810) South Block, Southampton General Hospital  
Tremona Road, Southampton, SO16 6YD, UK

**Development Phase:** Phase III

**Sponsor's Medical Monitor:** Marcin Mankowski

**Chief Investigator:** Professor Tom Wilkinson

**This document is the property of Synairgen Research Ltd and may not be used, divulged, published, or otherwise disclosed without written consent of Synairgen Research Ltd.**



**SYNAIRGEN RESEARCH LTD**

**STUDY PROTOCOL SG018 - PROTOCOL SIGNATURE PAGE**

**SPONSOR REPRESENTATIVE**

I have reviewed and agreed this protocol and its contents. My signature, in conjunction with the signatures of the Sponsor's Medical Officer, the Chief Investigator, confirms the agreement of all parties that the clinical trial will be conducted in accordance with the protocol and all applicable laws and regulations including, but not limited to, the International Council on Harmonisation Guideline for Good Clinical Practice (ICH GCP), the standards set out by the UK Policy Framework for Health and Social Care Research, The UK Medicines for Human Use (Clinical Trials) Regulations 2004 and associated amendments, the EU Clinical Trials Directive 2001/20/EC, the EU GCP Directive 2005/28/EC, the Clinical Trial Regulation (EU) No 536/2014, the US Federal Regulations, including but not limited to 21CFR and the Ethical principles that have their origin in the Declaration of Helsinki.

Signature:

DocuSigned by:  
  
 Signer Name: Richard Marsden  
Signing Reason: I approve this document  
Signing Time: 21 December 2020 | 6:36 AM PST  
9E3C95FB38A6499EB38C0E65BCAAC481

Name (PRINT):

Richard Marsden

Title:

CEO

Date of signature:

21 December 2020 | 6:36 AM PST

**This document is the property of Synairgen Research Ltd and may not be used, divulged, published, or otherwise disclosed without written consent of Synairgen Research Ltd.**


**SYNAIRGEN RESEARCH LTD**

**STUDY PROTOCOL SG018 - PROTOCOL SIGNATURE PAGE**

**SPONSOR'S MEDICAL OFFICER**

I have reviewed and agreed this protocol and its contents on behalf of the Sponsor. My signature, in conjunction with the signatures of the Sponsor Representative, the Chief Investigator, confirms the agreement of all parties that the research study will be conducted in accordance with the protocol and all applicable laws and regulations including, but not limited to, the International Council on Harmonisation Guideline for Good Clinical Practice (ICH GCP), the standards set out by the UK Policy Framework for Health and Social Care Research, The UK Medicines for Human Use (Clinical Trials) Regulations 2004 and associated amendments, the EU Clinical Trials Directive 2001/20/EC, the EU GCP Directive 2005/28/EC, the Clinical Trial Regulation (EU) No 536/2014, the US Federal Regulations, including but not limited to 21CFR and the Ethical principles that have their origin in the Declaration of Helsinki.

Signature:

DocuSigned by:  
*Marcin Mankowski*  
 Signer Name: Marcin Mankowski  
Signing Reason: I approve this document  
Signing Time: 21 December 2020 | 8:39 AM PST  
68CAB9DB86FC41A18CEF15571BD4C926

Marcin Mankowski

Name (PRINT):

Acting Chief Medical Officer

Title:

21 December 2020 | 8:39 AM PST

Date of signature:

**This document is the property of Synairgen Research Ltd and may not be used, divulged, published, or otherwise disclosed without written consent of Synairgen Research Ltd.**


**SYNAIRGEN RESEARCH LTD**

**CLINICAL STUDY PROTOCOL SG018 - PROTOCOL SIGNATURE PAGE**

**CHIEF INVESTIGATOR**

I have reviewed and agreed this protocol and its contents. My signature, in conjunction with the signatures of the Sponsor Representative, Sponsor's Medical Officer, confirms the agreement of all parties that the clinical trial will be conducted in accordance with the protocol and all applicable laws and regulations including, but not limited to, the International Council on Harmonisation Guideline for Good Clinical Practice (ICH GCP), the standards set out by the UK Policy Framework for Health and Social Care Research, The UK Medicines for Human Use (Clinical Trials) Regulations 2004 and associated amendments, the EU Clinical Trials Directive 2001/20/EC, the EU GCP Directive 2005/28/EC, the Clinical Trial Regulation (EU) No 536/2014, the US Federal Regulations, including but not limited to 21CFR and the Ethical principles that have their origin in the Declaration of Helsinki.

Signature:

DocuSigned by:  
*Tom Wilkinson*  
 Signer Name: Tom Wilkinson  
Signing Reason: I approve this document  
Signing Time: 21 December 2020 | 8:40 AM PST  
C61D5200A1B1471FA280272AF4BD82FB

Name (PRINT):

Tom wilkinson

Title:

Professor

Date of signature:

21 December 2020 | 8:40 AM PST

**This document is the property of Synairgen Research Ltd and may not be used, divulged, published, or otherwise disclosed without written consent of Synairgen Research Ltd.**

**SYNAIRGEN RESEARCH LTD**

**CLINICAL STUDY PROTOCOL SG018 - PROTOCOL SIGNATURE PAGE**

**PRINCIPAL INVESTIGATOR**

I confirm that I have read and agreed this protocol and its contents. I understand that all information concerning SNG001 and this protocol supplied to me by Synairgen Research Ltd is confidential.

My signature below confirms that as part of my responsibility as Principal Investigator, I will ensure that the clinical trial is conducted in accordance with the protocol and all applicable laws and regulations including, but not limited to, the International Council on Harmonisation Guideline for Good Clinical Practice (ICH GCP), the standards set out by the UK Policy Framework for Health and Social Care Research, The UK Medicines for Human Use (Clinical Trials) Regulations 2004 and associated amendments, the EU Clinical Trials Directive 2001/20/EC, the EU GCP Directive 2005/28/EC, the Clinical Trial Regulation (EU) No 536/2014, the US Federal Regulations, including but not limited to 21CFR and the Ethical principles that have their origin in the Declaration of Helsinki.

Signature:

Name (PRINT): \_\_\_\_\_

Title: \_\_\_\_\_

Date of signature: \_\_\_\_\_

**This document is the property of Synairgen Research Ltd and may not be used, divulged, published, or otherwise disclosed without written consent of Synairgen Research Ltd.**

## PROTOCOL SYNOPSIS

<b>Study Title:</b> A randomised, double-blind, placebo-controlled, Phase III trial to determine the efficacy and safety of inhaled SNG001 for the treatment of patients hospitalised due to moderate COVID-19	
<b>Protocol number:</b> SG018	
<b>Study centres:</b> Multi-national and multicentre	
<b>Number of randomised patients:</b> 610 randomised patients in total, randomised 1:1 to the following arms:  Arm 1: 305 patients - SNG001 - contents of 2 syringes* per dose Arm 2: 305 patients - Placebo - contents of 2 syringes <sup>+</sup> per dose  *Each syringe contains 0.65mL of SNG001 nebuliser solution, containing 12MIU/mL of interferon beta-1a <sup>+</sup> Each syringe contains 0.65mL of formulation buffer	
<b>Study period:</b> 2020 onwards	<b>Phase of development:</b> Phase III
<b>Background:</b> <p>Severe Acute Respiratory Syndrome coronavirus 2 (SARS-CoV-2) is a global threat and there is a need to assess new treatments which will prevent and effectively treat severe lower respiratory tract (LRT) illness caused by the SARS-CoV-2.</p> <p>Interferon beta (IFN-β) has showed antiviral activity against SARS-CoV-2 in cell-based assays (1). IFN-β driven anti-viral responses have been shown to be compromised/deficient in older people (2) and those with chronic airways diseases (3, 4). These, and other patient groups are at high risk of developing severe LRT illness which can be fatal (5). The IFN-β deficiency can be overcome through the administration of exogenous IFN-β. This has been shown both <i>in vitro</i>, using cells from patients, and in clinical trials using SNG001 (an inhaled IFN-β1a formulation for nebulisation). We hypothesise that SNG001 will rectify the deficiency in the lungs in at-risk patients and prevent severe LRT illness in the context of SARS-CoV-2 infection.</p> <p>The SG016 hospital pilot study was completed in May 2020. During this pilot study, 101 hospitalised adults, ≥18 years of age, with confirmed or suspected SARS-CoV-2 infection were randomised to receive SNG001 or placebo. Results of the pilot study showed that the risks of developing severe COVID-19 (the disease caused by SARS-CoV-2) were markedly reduced in patients receiving SNG001 compared to placebo and additionally that patients who received SNG001 were more than twice as likely to recover from COVID-19 as those on placebo. In addition, there was a significant reduction in breathlessness in patients receiving SNG001, compared to placebo (6).</p> <p>SNG001 has been well tolerated in all clinical studies to date. Around 280 patients have been treated with SNG001. Of the 280, approximately 50 had chronic obstructive pulmonary disease (COPD), 50 had confirmed COVID-19 with varying underlying diseases i.e. heart</p>	

disease, lung disease, diabetes etc and the remaining 180 had asthma. The majority of the 280 patients had or were suspected to have an active respiratory viral infection (rhinovirus, influenza, coronavirus, SARS-CoV-2, etc) at the time of randomisation.

SNG001 is pH neutral, rather than acidic and does not contain excipients such as mannitol, human serum albumin (HSA) and arginine, which are present in the injectable IFN- $\beta$  formulations and which may have their own unwanted effects if delivered to the lungs.

SNG001 has historically been delivered using the I-neb, a mesh nebuliser made by Philips Respironics. The I-neb has been tested to ensure the drug retains its activity after aerosolization. A dose escalating trial established a target lung dose which induced an antiviral response in the lungs that was present 24 hours after dose administration.

In this trial the Aerogen Ultra device will be used. The Ultra is mesh nebuliser that is widely available and is better suited to single patient usage in the hospital setting. Laboratory assessments found that both the I-neb and the Ultra had similar levels of protein content in and similar IFN- $\beta$  activity post nebulisation.

The primary endpoint is recovery in patients with confirmed SARS-CoV-2 infection who are hospitalised due to moderate COVID-19, after administration of SNG001 compared to placebo, where moderate COVID-19 is defined as presence of clinical signs and symptoms necessitating administration of oxygen therapy by mask or nasal prongs and recovery is defined as no limitation of activities according to the Ordinal Scale of Clinical Improvement (OSCI), with no rebound at subsequent assessments. The OSCI to be used in this trial is the 18<sup>th</sup> February 2020 version as recommended by the World Health Organization (WHO) (7).

**Purpose of the study:** The purpose of this Phase III study is to confirm that SNG001 can accelerate the recovery of hospitalised patients receiving oxygen with confirmed SARS-CoV-2. Safety and other efficacy endpoints will also be assessed.

**Study Population:**

Adults,  $\geq 18$  years of age with SARS-CoV-2 infection confirmed by a positive virus test using a validated molecular assay e.g. Reverse Transcription Polymerase Chain Reaction (RT-PCR) or validated antigen assay e.g. Sofia 2 SARS Antigen FIA, who are hospitalised due to COVID-19 and require oxygen therapy, either via nasal prongs or a mask, but do not require high-flow oxygen therapy, continuous positive airway pressure (CPAP) treatment or intubation.

**Study Design:**

Eligible patients will be randomised in a 1:1 ratio to receive SNG001 two syringes or placebo two syringes.

Patients who had positive virus test for SARS-CoV-2 prior to hospitalisation will be randomised no later than 48 hours after hospital admission. If the virus test was performed more than 96 hours prior to hospitalisation, the test will have to be repeated in the hospital prior to randomisation. Only patients whose repeated virus test is positive will be randomised, no later than 48 hours after confirmation of SARS-CoV-2 infection.

Patients who had positive virus test for SARS-CoV-2 after hospitalisation will be randomised no later than 48 hours after confirmation of SARS-CoV-2 infection.

SNG001 or placebo will be administered via the Ultra nebuliser. Patients will receive a dose

of SNG001 or placebo once a day for 14 days and will be followed up for up to 90 days after completion of study medication. Study data will be collected from patients daily, as per the study schedule. Efficacy will be determined through differences between the groups in the OSCI scores, and the secondary endpoints. Adverse events (AEs) and concomitant medications will be monitored throughout the study period.

A Data Safety Monitoring Committee (DSMC) will perform a review of the safety data before 100 patients complete study treatment, to ensure the safety of study patients. The DSMC will also meet as and when necessary, i.e. if a safety issue arises or when the DSMC requests a further meeting.

### **Study Objectives**

#### **Primary Objective:**

To evaluate recovery in patients with moderate COVID-19 after administration of SNG001 compared to placebo.

#### **Secondary Objectives:**

- a. To evaluate the efficacy of SNG001 compared to placebo in patients with moderate COVID-19, using a range of endpoints.
- b. To assess the general safety and tolerability of SNG001 compared to placebo when administered to patients with moderate COVID-19.

### **Study Endpoints**

#### **Primary Endpoints:**

- a. Time to hospital discharge, defined by the OSCI score of 2 or below, with no rebound at subsequent assessments.
- b. Time to recovery, where recovery is defined as the OSCI score of 1 or below, with no rebound at subsequent assessments.

#### **Key Secondary Endpoints:**

- a. Progression to severe disease or death, defined by the OSCI score of 5 or above within 35 days of first dose.
- b. Progression to intubation or death, defined by the OSCI score of 6 or above within 35 days of first dose.
- c. Death within 35 days of first dose.

#### **Secondary Endpoints:**

- a. Recovery, where recovery is defined as the OSCI score of 1 or below, with no rebound at subsequent assessments, at Days 7, 14, 21 and 28.
- b. Hospital discharge at Days 7, 14, 21 and 28.
- c. Improvement across the entire OSCI at Days 7, 14, 21 and 28.
- d. Changes in breathlessness, cough and sputum scale (BCSS) score during the study period, including disaggregated breathlessness and cough scores.
- e. Changes in National Early Warning Score 2 (NEWS2) during the hospitalisation period.
- f. Daily assessment of COVID-19 symptoms and limitation of usual activities.
- g. Quality of life measured using EQ-5D-5L.
- h. Long-COVID-19 symptoms.
- i. Safety and tolerability – vital signs, AEs and concomitant medications.

### **Inclusion criteria:**

To be eligible for randomisation into this study, each patient must fulfil the following criteria:

1. Male or female,  $\geq 18$  years of age at the time of consent.
2. Admitted to hospital due to the severity of their COVID-19.
3. Positive virus test for SARS-CoV-2 using a validated molecular assay or antigen assay. Patients who had positive virus test for SARS-CoV-2 prior to hospitalisation will be randomised no later than 48 hours after hospital admission. If the virus test was performed more than 96 hours prior to hospitalisation, the test will have to be repeated in the hospital prior to randomisation. Only patients whose repeated virus test is positive will be randomised, no later than 48 hours after confirmation of SARS-CoV-2 infection.

Patients who had positive virus test for SARS-CoV-2 after hospitalisation will be randomised no later than 48 hours after confirmation of SARS-CoV-2 infection.

4. Require oxygen therapy via nasal prongs or mask (OSCI score of 4).
5. Provided informed consent.
6. Female patients must be  $\geq 1$  year post-menopausal, surgically sterile, or using a highly effective method of contraception. Acceptable highly effective methods of contraception include;
  - bilateral tubal occlusion
  - intrauterine device (provided coils are copper-banded)
  - levonorgestrel intrauterine system (e.g., Mirena™)
  - medroxyprogesterone injections (e.g., Depo-Provera™)
  - etonogestrel implants (e.g., Implanon™, Norplan™)
  - normal and low dose combined oral pills
  - norelgestromin/ ethinylestradiol transdermal system
  - intravaginal device (e.g., ethinylestradiol and etonogestrel), desogestrel (e.g., Cerazette™)
  - total sexual abstinence (defined as refraining from heterosexual intercourse)
  - vasectomised sexual partner.

Women should have been stable on their chosen method of birth control for a minimum of 3 months before entering the trial and should continue with birth control for 1 month after the last dose of inhaled IFN- $\beta$ 1a/matching placebo. In addition to the highly effective method of contraception (except for the practice of total sexual abstinence), a condom (in UK with spermicides) should be used by the male partner for sexual intercourse from randomisation (Visit 2) and for 1 month after the last dose of inhaled IFN- $\beta$ 1a/matching placebo to prevent pregnancy.

7. Women not of childbearing potential are defined as women who are either permanently sterilized (hysterectomy, bilateral oophorectomy, or bilateral salpingectomy), or who are postmenopausal. Women will be considered post-menopausal if they have been amenorrhoeic for 12 months prior to the planned date of randomisation without an alternative medical cause. The following age specific requirements apply:
  - Women  $< 50$  years old would be considered post-menopausal if they have been amenorrhoeic for 12 months or more following cessation of exogenous

hormonal treatment and if follicle stimulating hormone (FSH) levels are in the postmenopausal range.

- Women  $\geq 50$  years old would be considered post-menopausal if they have been amenorrhoeic for 12 months or more following cessation of all exogenous hormonal treatment.

If, in the setting of the pandemic, the use of an acceptable birth control method is not possible, the decision to enrol a woman of childbearing potential should be based on the benefit-risk for the patient, which should be discussed with the patient at the time of the informed consent.

### **Exclusion Criteria:**

A patient must not be randomised into the study if they meet any of the following criteria:

1. Evidence of ongoing SARS-CoV-2 infection for more than 3 weeks, confirmed by a validated molecular assay or validated antigen assay.
2. Non-invasive ventilation or high-flow oxygen (OSCI score of 5).
3. Mechanical ventilation (continuous or intermittent CPAP or intubation) or admission to intensive care (OSCI score of  $\geq 6$ ).
4. Previous SARS-CoV-2 infection confirmed by a validated molecular assay or validated antigen assay.
5. Any condition, including findings in the patients' medical history or in the pre-randomisation study assessments that in the opinion of the Investigator, constitute a risk or a contraindication for the participation of the patient into the study or that could interfere with the study objectives, conduct or evaluation.
6. Participation in previous clinical trials of SNG001.
7. Current or previous participation in another clinical trial where the patient has received a dose of an Investigational Medicinal Product (IMP) containing small molecules within 30 days or 5 half-lives (whichever is longer) prior to entry into this study or containing biologicals within 3 months prior to entry into this study.
8. Inability to use a nebuliser with a mouthpiece.
9. Inability to comply with the requirements for storage conditions of study medication in the home setting.
10. History of hypersensitivity to natural or recombinant IFN- $\beta$  or to any of the excipients in the drug preparation.
11. Females who are breast-feeding, lactating, pregnant or intending to become pregnant.
12. Previous SARS-CoV-2 vaccination.

**Test product, dose, and mode of administration:** SNG001 nebuliser solution is presented in glass syringes containing 0.65 ml of drug product solution at a concentration of 12 MIU/ml.

The Ultra device will be filled with the contents of two syringes (if randomised to Arm 1). Patients will inhale study medication once a day.

**Reference product, dose and mode of administration:** The placebo will be the same formulation as the study medication but without IFN- $\beta$ 1a (i.e. only the excipients of the SNG001 solution).

The Ultra device will be filled with the contents of two syringes (if randomised to Arm 2). Patients will inhale study medication once a day.

**Duration of Treatment:** Patients will receive study medication for 14 days.

**Sample Size:** Success will be determined if at least one of the primary endpoints is declared statistically significant by the primary analysis. A sample size of 610 patients in total using a 1:1 randomisation ratio (305 patients per treatment arm) has been chosen to provide at least 90% power to detect a hazard ratio of 1.45 in time to hospital discharge and a hazard ratio of 1.7 in time to recovery and at least 95% power to declare statistical significance on at least one of the primary endpoints. This sample size has been calculated using a global 2-sided alpha level of 0.05 and allows for an interim analysis to assess futility.

## **Statistical Methods**

### **Analysis populations**

The intention-to-treat (ITT) population is defined as all randomised patients.

The safety population will include all randomised patients who receive at least one dose of study drug.

The per protocol population will include all patients in the ITT population who do not have any protocol deviations with an impact on efficacy on or prior to Day 35.

### **Statistical analysis methodology**

The same method of analysis will be used for both primary endpoints. The primary analysis will use a Cox proportional hazards model. Patients who die will be administratively censored at Day 28, i.e. time to hospital discharge and time to recovery for these patients will be fixed at 28 days. Patients who neither recover nor die will be censored at the date of their last OSCI assessment.

For the key secondary endpoint (progression to severe disease or death) a logistic regression model will be used to estimate the difference in probabilities between the treatment groups.

Safety analyses will be based primarily on AE information which will be summarized descriptively, including summary of treatment-emergent AEs.

## **Estimands**

### **Primary Estimand**

The first primary estimand will be the hazard ratio of SNG001 versus placebo for the time to hospital discharge of patients in the ITT population. Intercurrent events will be handled as per the table below.

<b>Intercurrent event</b>	<b>Policy</b>
Death	Composite variable: Patients who die on or prior to day 28 will have time to hospital discharge right censored at 28 days (the maximum time to discharge).

Premature treatment discontinuation and missed doses	Treatment policy: The time to hospital discharge will be used regardless of treatment discontinuation and the number of missed doses.
Withdrawal from study following hospital discharge	Treatment policy: The time to hospital discharge will be used regardless of study withdrawal and regardless of the duration of follow-up following hospital discharge. This assumes relapse is independent to the treatment effect.
Withdrawal from study prior to hospital discharge	While-on-study: Time to hospital discharge will be right censored at the time of withdrawal from the study or at 28 days, whichever occurs first. The evaluation period for this endpoint is 28 days, therefore estimating the time to hospital discharge after Day 28 would not be appropriate. Premature study withdrawal is expected to be independent to the treatment effect, therefore a censoring approach in this scenario is considered appropriate. However, this assumption will be tested by appropriate sensitivity analyses.
Other protocol deviations and concomitant medications affecting efficacy	Treatment policy: Time to hospital discharge will be used regardless of any concomitant medication use or protocol deviations.

The second primary estimand will be the hazard ratio of SNG001 versus placebo for the time to recovery of patients in the ITT population. Intercurrent events will be handled as per the table below.

Intercurrent event	Policy
Death	Composite variable: Patients who die during the study period will have time to recovery right censored at 28 days (the maximum possible time to recovery).
Premature treatment discontinuation and missed doses	Treatment policy: The time to recovery will be used regardless of treatment discontinuation and the number of missed doses.
Withdrawal from study following recovery	Treatment policy: The time to recovery will be used regardless of study withdrawal and regardless of the duration of follow-up following recovery. This assumes relapse is independent to the treatment effect.

Withdrawal from study prior to recovery	While-on-study: Time to recovery will be right censored at the time of withdrawal from the study or at 28 days, whichever occurs first. The evaluation period for this endpoint is 28 days, therefore estimating the time to recovery after Day 28 would not be appropriate. Premature study withdrawal is expected to be independent to the treatment effect, therefore a censoring approach in this scenario is considered appropriate. However, this assumption will be tested by appropriate sensitivity analyses.										
Other protocol deviations and concomitant medications affecting efficacy	Treatment policy: Time to recovery will be used regardless of any concomitant medication use or protocol deviations.										
<p><b>Key Secondary Estimands</b></p> <p>difference in probabilities for SNG001 versus placebo for progression to severe disease or death within 35 days of first dose of patients in the ITT population. Intercurrent events will be handled as follows:</p> <table> <tr> <th>Intercurrent event</th><th>Policy</th></tr> <tr> <td>Death</td><td>Composite variable: The definition of severe disease includes patients who die within 35 days of first dose, regardless of their OSCI assessments.</td></tr> <tr> <td>Premature treatment discontinuation and missed doses</td><td>Treatment policy: Severe disease will be determined regardless of treatment discontinuation and the number of missed doses.</td></tr> <tr> <td>Withdrawal from study</td><td>While-on-study: Patients will only be considered having severe disease if observed and recorded whilst participating in the study. This assumes the treatment effect is independent to reason for study withdrawal.</td></tr> <tr> <td>Other protocol deviations and concomitant medications affecting efficacy</td><td>Treatment policy: Severe disease will be determined regardless of any concomitant medication use or protocol deviations.</td></tr> </table> <p>The second key secondary estimand will be the difference in probabilities for SNG001 versus placebo for progression to intubation or death within 35 days of first dose of patients in the ITT population.</p>		Intercurrent event	Policy	Death	Composite variable: The definition of severe disease includes patients who die within 35 days of first dose, regardless of their OSCI assessments.	Premature treatment discontinuation and missed doses	Treatment policy: Severe disease will be determined regardless of treatment discontinuation and the number of missed doses.	Withdrawal from study	While-on-study: Patients will only be considered having severe disease if observed and recorded whilst participating in the study. This assumes the treatment effect is independent to reason for study withdrawal.	Other protocol deviations and concomitant medications affecting efficacy	Treatment policy: Severe disease will be determined regardless of any concomitant medication use or protocol deviations.
Intercurrent event	Policy										
Death	Composite variable: The definition of severe disease includes patients who die within 35 days of first dose, regardless of their OSCI assessments.										
Premature treatment discontinuation and missed doses	Treatment policy: Severe disease will be determined regardless of treatment discontinuation and the number of missed doses.										
Withdrawal from study	While-on-study: Patients will only be considered having severe disease if observed and recorded whilst participating in the study. This assumes the treatment effect is independent to reason for study withdrawal.										
Other protocol deviations and concomitant medications affecting efficacy	Treatment policy: Severe disease will be determined regardless of any concomitant medication use or protocol deviations.										

The third key secondary estimand will be the difference in probabilities for SNG001 versus placebo for death within 35 days of first dose of patients in the ITT population. For both the second and third key secondary endpoints the same policies will be used for handling intercurrent events as progression to severe disease or death.

### **Interim Analyses**

An unblinded interim analysis is planned to assess futility on the first 300 randomised patients and will be conducted once all of these patients have completed the day 35 visit or have withdrawn from the study. The primary analyses for the two primary endpoints will be conducted for the interim analysis and provided to an Independent Data Monitoring Committee (IDMC).

## Study Schedule

Assessment	Pre-Treatment <sup>1</sup>	Treatment Phase <sup>2</sup>	End of Treatment <sup>3</sup>	Follow-up		
	Day 0	Days 1 to 14	Day 15 (+1 day)	Day 16 to Day 28	Day 29 to Day 35	Day 36 to Day 90
Informed consent	X					
Inclusion / Exclusion criteria	X					
Medical history	X					
COVID 19 infection history	X					
Patient demographics	X					
Physical examination	X <sup>4</sup>	X <sup>4,5</sup>	X <sup>4,5</sup>			
Height and weight	X					
Ordinal Scale of Clinical Improvement	X	X <sup>6,7</sup>	X <sup>6,7</sup>	X <sup>6,7</sup>	X <sup>6,7</sup>	
BCSS	X	X <sup>6,7</sup>	X <sup>6,7</sup>	X <sup>6,7</sup>		X <sup>6,7,8</sup>
Vital signs <sup>9</sup>	X	X <sup>5</sup>	X <sup>5</sup>			
NEWS2	X	X <sup>5</sup>	X <sup>5</sup>	X <sup>5</sup>		
EQ-5D-5L	X <sup>10</sup>	X <sup>6,7,11</sup>	X <sup>6,7,10</sup>	X <sup>6,7,11</sup>	X <sup>6,7,11</sup>	X <sup>6,7,8</sup>
COVID-19 symptom assessment		X <sup>6,7</sup>	X <sup>6,7</sup>	X <sup>6,7</sup>		
Long-COVID-19 assessments			X <sup>6,7</sup>	X <sup>6,7,12</sup>		X <sup>6,7,8</sup>
Urine pregnancy test (if applicable)	X					
Blood samples for immunogenicity test <sup>13</sup>	X					X <sup>14</sup>
Device training (if required)	X	X				
Randomisation	X					
Dose administration		X				
Recording of AEs and SAEs <sup>15</sup>	X	X <sup>6,7</sup>	X <sup>6,7</sup>	X <sup>6,7</sup>	X <sup>6,7</sup>	X <sup>6,7</sup>
Recording of concomitant medications	X	X <sup>6,7</sup>	X <sup>6,7</sup>	X <sup>6,7</sup>	X <sup>6,7</sup>	X <sup>6,7</sup>

1. Dosing may start on the same day as the pre-treatment assessments and if so the Day 1 assessments will not be performed
2. All assessments and sampling are to be carried out pre-dose
3. If Day 15 is +1 day, day 16 will be missed and no deviation recorded
4. Includes chest exam only unless a physical examination is deemed necessary by the Investigator
5. Hospitalised patients only
6. For patients who have been discharged, assessments should be performed via phone/video link, if possible
7. Patients who stop dosing (for example if they become ventilated or for any other reason) should undergo these key data assessments if possible
8. Day 60 and Day 90 (+/- 2 days) only
9. Includes daily temperature, respiratory rate, heart rate and blood pressure
10. Complete EQ-5D-5L questionnaire
11. Complete EQ-5D-5L questionnaire on Day 7 and Day 28 only. The mobility, self-care and usual activities dimensions of the questionnaire will be assessed daily from Day 1 to Day 35
12. Day 28 only
13. All sites/patients, where collection is feasible
14. Day 44 (+/-5 days) only
15. See section 7.7.1 and 7.8 for AE and SAE reporting requirements

## TABLE OF CONTENTS

PROTOCOL SYNOPSIS .....	7
TABLE OF CONTENTS.....	17
1 INTRODUCTION.....	23
2 STUDY OBJECTIVES AND ENDPOINTS .....	24
2.1 Objectives .....	24
2.2 Primary Endpoints:.....	24
2.3 Key Secondary Endpoint:.....	25
2.4 Secondary Endpoints:.....	25
3 STUDY DESIGN .....	25
3.1 Criteria for Inclusion and Exclusion .....	26
3.1.1 Inclusion Criteria.....	26
3.1.2 Exclusion Criteria.....	27
4 STUDY MEDICATION .....	28
4.1 Allocation to Study Medication .....	28
4.2 Breaking the Blind.....	28
4.3 Stopping Rules .....	28
4.3.1 Individual Patient Stopping Rules .....	28
4.3.2 Trial Stopping Rule .....	29
4.4 Description of Investigational Medicinal Product and Device .....	29
4.4.1 SNG001 .....	29
4.4.2 Placebo .....	29
4.4.3 Inhalation Device .....	30
4.5 Dosage and Administration .....	30
4.5.1 Dose Delivery Times.....	30
4.5.2 Packaging and Labelling .....	30
4.6 Preparation, Handling and Dispensing .....	31
4.6.1 Preparation Time .....	31
4.6.2 Timing of Doses .....	31
4.6.3 Missed or Incomplete Doses .....	31
4.7 Storage of Study Medication.....	31
4.8 Drug Accountability .....	31
4.9 Treatment of Overdose .....	32
5 STUDY PROCEDURES.....	33
5.1 Patient Numbering.....	33
5.2 Screening Failures .....	33
5.3 Study Visits .....	33
5.3.1 Pre-Treatment.....	33

5.3.2	Days 1 to 14.....	34
5.3.3	End of Treatment.....	35
5.3.4	Follow-up .....	36
5.4	Unscheduled Visits.....	36
5.5	Patient Withdrawal .....	37
6	STUDY ASSESSMENTS.....	37
6.1	Informed Consent .....	37
6.2	Medical History .....	37
6.3	COVID-19 Infection History.....	38
6.4	Patient Demography .....	38
6.5	Physical examination.....	38
6.6	Ordinal Scale for Clinical Improvement .....	39
6.7	Breathlessness, Cough and Sputum Score (BCSS).....	40
6.8	Vital Signs .....	41
6.9	National Early Warning Score 2 (NEWS2) .....	41
6.10	Quality of life measured using EQ-5D-5L.....	41
6.11	COVID-19 symptom assessment .....	42
6.12	Assessment of Long-COVID-19 symptoms.....	42
6.13	Samples for Laboratory Assessment .....	42
6.13.1	Urine Pregnancy Test .....	42
6.13.2	Blood sample for immunogenicity testing .....	43
6.14	Concomitant Medications.....	43
6.15	Prohibited Medications.....	43
7	ADVERSE EVENT REPORTING .....	44
7.1	Adverse Events.....	44
7.2	Treatment-Emergent Adverse Events .....	44
7.3	Serious Adverse Event .....	45
7.4	Assessment of Adverse Event .....	45
7.4.1	Relationship to Study Medication .....	45
7.5	Discontinuation of Study Medication due to an AE.....	47
7.6	Pregnancy .....	47
7.7	Adverse Events Recording and Reporting .....	47
7.7.1	Reporting of Adverse Events .....	47
7.7.2	Recording of Adverse Events.....	47
7.8	Reporting of Serious Adverse Events .....	48
8	DATA ANALYSIS/STATISTICAL METHODS .....	49
8.1	Sample Size Determination .....	49
8.2	Randomisation.....	50
8.3	Analysis Populations .....	50

8.4	Estimands and Intercurrent Events.....	50
8.4.1	Primary Efficacy Estimands.....	50
8.4.2	Key Secondary Efficacy Estimands .....	54
8.5	Statistical Analysis Methodology for Efficacy Endpoints .....	55
8.5.1	Analysis of the Primary Efficacy Endpoint.....	55
8.5.2	Analysis of the Key Secondary Efficacy Endpoints .....	56
8.5.3	Analysis of Secondary Efficacy Endpoints .....	56
8.5.4	Multiple Testing Strategy .....	57
8.5.5	Handling of Missing Data .....	57
8.6	Safety Analysis.....	57
8.7	Other Analyses .....	58
8.8	Subgroup Analyses.....	58
8.9	Interim Analyses.....	58
8.10	Primary Analysis .....	59
8.11	Data Safety Monitoring Committee .....	59
9	QUALITY CONTROL AND QUALITY ASSURANCE.....	59
9.1	Quality Control (Monitoring).....	59
9.2	Quality Assurance .....	60
10	DATA HANDLING AND RECORD KEEPING.....	60
10.1	Study Source Documents and Electronic Case Report Form.....	60
10.2	Record Retention.....	61
11	REGULATORY AND ETHICS .....	61
11.1	Regulatory and Ethics Considerations .....	61
11.2	Competent Authority Approval.....	61
11.3	Institutional Review Board/Independent Ethics Committee and Hospital Board Requirements.....	61
11.4	Informed Consent and Screening Data.....	62
11.5	Patient Confidentiality.....	62
11.6	Investigator Compliance.....	62
11.7	Progress Reports and Safety Reports .....	63
12	DEFINITION OF END OF TRIAL .....	63
13	SPONSOR DISCONTINUATION CRITERIA.....	63
13.1	Study Sites.....	63
13.2	Entire Study .....	63
14	PUBLICATION OF STUDY RESULTS.....	63
15	APPENDICES.....	65
	<b>APPENDIX A: AEROGEN ULTRA DEVICE.....</b>	<b>65</b>
	<b>APPENDIX B: WITHDRAWAL FLOW CHART.....</b>	<b>69</b>
	<b>APPENDIX C: BCSS QUESTIONNAIRE.....</b>	<b>70</b>

APPENDIX D: EQ-5D-5L .....	71
APPENDIX E: LONG-COVID-19 PATIENT REPORTED OUTCOME MEASURES .....	73
16 REFERENCES .....	78

## ABBREVIATIONS AND DEFINITIONS

<b>Abbreviation/Terms</b>	<b>Definition</b>
AE	Adverse Event
BCSS	Breathlessness, Cough and Sputum Scale
BMI	Body Mass Index
CI	Confidence Interval
COPD	Chronic Obstructive Pulmonary Disease
COVID-19	Disease caused by SARS-CoV-2
CPAP	Continuous Positive Airway Pressure
CRA	Clinical Research Associate
CRO	Clinical Research Organisation
Delegate	A qualified member of the study team delegated to perform a task by the investigator
DSMC	Data Safety Monitoring Committee
eCRF	Electronic Case Report Form
EQ-5D-5L	EuroQuol 5-dimension 5-level (quality of life questionnaire)
FDA	U.S. Food and Drug Administration
FEV <sub>1</sub>	Forced Expiratory Volume in 1 second
FSH	Follicle Stimulating Hormone
GCP	Good Clinical Practice
GLP	Good Laboratory Practice
GMP	Good Manufacturing Practice
HSA	Human Serum Albumin
IB	Investigator's Brochure
ICH	International Council for Harmonisation
IDMC	Independent Data Monitoring Committee
IEC	Independent Ethics Committee
IFN-β	Interferon-beta
IMP	Investigational Medicinal Product
IRB	Institutional Review Board
ITT	Intention-to-Treat
LRT	Lower Respiratory Tract
MedDRA	Medical Dictionary for Regulatory Activities
MHRA	Medicines and Healthcare products Regulatory Agency
MIU	Million International Units
MMRM	Mixed Models with Repeated Measures
NEWS2	National Early Warning Score (NEWS) 2
NOAEL	No-observed-adverse-effect-level
Placebo	Comparator Medication Without the Active Ingredient
R&D	Research and Development
Randomisation	Random Allocation to Treatment Group
REC	Research Ethics Committee
RTI	Respiratory Tract Infection
RT-PCR	Reverse Transcription Polymerase Chain Reaction
SAE	Serious Adverse Event
SAP	Statistical Analysis Plan

SARS-CoV-2	Severe Acute Respiratory Syndrome-Coronavirus 2
SOC	Standard of Care
Source data	Original documents, data and record (including but not limited to: hospital records, study source documents, lab reports)
Sponsor	Company that takes responsibility for the initiation, management and financing of the trial: Synairgen Research Limited
TEAE	Treatment-Emergent Adverse Event
VAS	Visual Analogue Scale

## 1 INTRODUCTION

Severe Acute Respiratory Syndrome coronavirus 2 (SARS-CoV-2) is a global threat and there is a need to assess new treatments which will prevent and effectively treat severe lower respiratory tract (LRT) illness caused by the SARS-CoV-2 virus.

Interferon beta (IFN-β) has showed antiviral activity against SARS-CoV-2 in cell-based assays (1). IFN-β driven anti-viral responses have been shown to be compromised/deficient in older people (2) and those with chronic airways diseases (3, 4). These, and other patient groups are at high risk of developing severe LRT illness which can be fatal (5). The IFN-β deficiency can be overcome or boosted through the administration of exogenous IFN-β. This has been shown both *in vitro*, using cells from patients, and in clinical trials using SNG001 (an inhaled IFN-β1a formulation for nebulisation). We hypothesise that SNG001 will rectify the deficiency in the lungs in at-risk patients and prevent severe LRT illness in the context of SARS-CoV-2 infection.

The SG016 hospital pilot study was completed in May 2020. During this pilot study, 101 hospitalised adults, ≥18 years of age, with confirmed or suspected SARS-CoV-2 infection were randomised to receive SNG001 or placebo. Results of the pilot study showed that the risks of developing severe COVID-19 (the disease caused by SARS-CoV-2) were markedly reduced in patients receiving SNG001 compared to those receiving placebo and that patients who received SNG001 were more than twice as likely to recover from COVID-19 as those on placebo. In addition, there was a significant reduction in breathlessness in patients receiving SNG001, compared to placebo (6).

SNG001 has been well tolerated in all clinical studies to date. Around 280 patients have been treated with SNG001. Of these, approximately 50 had chronic obstructive pulmonary disease (COPD), 50 had confirmed COVID-19 with varying underlying diseases i.e. heart disease, lung disease, diabetes etc and the remaining 180 had asthma. The majority of the 280 patients had or were suspected to have an active respiratory viral infection (rhinovirus, influenza, coronavirus, SARS-CoV-2, etc) at the time of randomisation.

SNG001 is pH neutral, rather than acidic and does not contain excipients such as mannitol, human serum albumin (HSA) and arginine, which are present in the injectable IFN-β formulations and which may have their own unwanted effects if delivered to the lungs.

SNG001 has historically been delivered using the I-neb, a mesh nebuliser made by Philips Respironics. The I-neb has been tested to ensure the drug retains its activity after aerosolization. A dose escalating trial established a target lung dose which induced an antiviral response in the lungs that was present 24 hours after dose administration.

This SG018 trial is designed to assess the efficacy of a 2 syringe dose of SNG001 delivered by the Ultra nebulizer. Two syringes in the Ultra are predicted to deliver a lung dose of approximately 5 MIU, which is approximately 32% higher than has historically been delivered by the I-neb (3.8 MIU).

The rationale for the slightly higher dose administered with the Ultra nebulizer is based on the non-clinical data that support a maximum administered lung dose per day to humans of 132 µg (33 MIU), assuming a conservative 60 kg human body weight. This dose is based on the no-observed-adverse-effect-level (NOAEL) from the Good Laboratory Practice (GLP) repeat dose 14-day toxicity study in cynomolgus monkey of 8.7 µg/kg/day total inhaled dose,

which corresponds to 2.2 µg/kg/day lung deposited dose, assuming a lung deposition fraction of 25%. The maximum dose that will be administered to patients in this study from the two syringes of SNG001 will be 57.1 µg (14.3 MIU) per day. The dose of 57.1 µg (14.3 MIU) represents the emitted dose from the Ultra nebulizer and therefore, the proportion of drug delivered to the lung will be lower. The estimated administered lung dose with the Ultra nebulizer will be 34.9% of the emitted dose of 57.1 µg (14.3 MIU) and this equates to a lung dose of 20 µg (5 MIU) (9). Safety margins of at least 4.2-fold between the NOAEL in the GLP toxicity study in monkeys and the dose of 57.1 µg/day (14.3 MIU) administered in this study have been calculated, based on comparative lung exposure (mg/g lung) and using the U.S. Food and Drug Administration (FDA)-accepted method by Tepper et al. (10).

Safety of SNG001 has been investigated in 5 completed clinical studies, across a variety of patient populations including patients with asthma, patients with COPD and patients hospitalized due to severity of COVID-19. Treatment has been well tolerated, and no specific safety signals were observed, whether related to local tolerance or systemic effects. Based on the extensive experience from parenteral IFN-β1a administration, and negligible systemic exposure after administration of SNG001, clinically significant systemic events are not anticipated. The data from pre-clinical studies and the previous clinical trials with SNG001 do not indicate risk for local adverse effects in the lungs at the proposed doses.

The primary endpoint of this study is to evaluate recovery in hospitalised patients receiving oxygen with confirmed SARS-CoV-2 infection after administration of SNG001 compared to placebo, where recovery is defined as the Ordinal Scale of Clinical Improvement (OSCI) score of 1 or below, with no rebound at subsequent assessments. The OSCI to be used in this trial is the 18th February 2020 version as recommended by the World Health Organisation (WHO) (7).

Complete information on SNG001 is available in the Investigator's Brochure (IB).

## **2 STUDY OBJECTIVES AND ENDPOINTS**

### **2.1 Objectives**

Primary objective:

To evaluate recovery in patients with moderate COVID-19 after administration of SNG001 compared to placebo.

Secondary objectives:

- a. To evaluate the efficacy of SNG001 compared to placebo in patients with moderate COVID-19, using a range of endpoints.
- b. To assess the general safety and tolerability of SNG001 compared to placebo when administered to patients with moderate COVID-19.

### **2.2 Primary Endpoints:**

- a. Time to hospital discharge, defined by the OSCI score of 2 or below, with no rebound at subsequent assessments.
- b. Time to recovery, where recovery is defined as the OSCI score of 1 or below, with no

rebound at subsequent assessments.

### **2.3 Key Secondary Endpoint:**

- a. Progression to severe disease or death, defined by the OSCI score of 5 or above within 35 days of first dose.
- b. Progression to intubation or death, defined by the OSCI score of 6 or above within 35 days of first dose.
- c. Death within 35 days of first dose.

### **2.4 Secondary Endpoints:**

- a. Recovery, where recovery is defined as the OSCI score of 1 or below, with no rebound at subsequent assessments, at days 7, 14, 21 and 28.
- b. Hospital discharge at days 7, 14, 21 and 28.
- c. Improvement across the entire OSCI at days 7, 14, 21 and 28.
- d. Changes in breathlessness, cough and sputum scale (BCSS) score during the study period, including disaggregated breathlessness and cough scores.
- e. Changes in National Early Warning Score (NEWS)2 during the hospitalisation period.
- f. Daily assessment of COVID-19 symptoms and limitation of usual activities.
- g. Quality of life measured using EuroQol 5-dimension 5-level (EQ-5D-5L).
- h. Long-COVID-19 symptoms.
- i. Safety and tolerability – vital signs, adverse events (AEs) and concomitant medications.

## **3 STUDY DESIGN**

610 adults,  $\geq 18$  years of age with SARS-CoV-2 infection confirmed by a positive virus test using a validated molecular assay e.g. Reverse Transcription Polymerase Chain Reaction (RT-PCR) or antigen assay, e.g. Sofia 2 SARS Antigen FIA, who are hospitalised due to COVID-19 and require oxygen therapy, either via nasal prongs or a mask, but do not require high-flow oxygen therapy, continuous positive airway pressure (CPAP) treatment or intubation will be randomised 1:1 to receive SNG001 two syringes (305 patients) or placebo two syringes (305 patients).

Patients who had positive virus test for SARS-CoV-2 prior to hospitalisation will be randomised no later than 48 hours after hospital admission. If the virus test was performed more than 96 hours prior to hospitalisation, the test will have to be repeated in the hospital prior to randomisation. Only patients whose repeated virus test is positive will be randomised, no later than 48 hours after confirmation of SARS-CoV-2 infection.

Patients who had positive virus test for SARS-CoV-2 after hospitalisation will be randomised no later than 48 hours after confirmation of SARS-CoV-2 infection.

SNG001 or placebo will be administered via the Ultra nebuliser. Patients will receive a dose of SNG001 (two syringes) or placebo (two syringes) once a day for 14 days and will be followed for up to 90 days after completion of study medication. Study data will be collected from patients as per the study schedule. Efficacy will be determined though differences between the groups in the time to recovery (primary endpoint), time to hospital discharge and

progression to severe disease or death (key secondary endpoints), and the secondary endpoints. AEs and concomitant medications will be monitored throughout the study period.

To ensure patient safety, a Data Safety Monitoring Committee (DSMC) will perform a review of the safety data before 100 patients complete study treatment, to ensure safety of study patients. The DSMC will also meet as and when necessary i.e. if a safety issue arises or when the DSMC requests a further meeting.

### **3.1 Criteria for Inclusion and Exclusion**

#### **3.1.1 Inclusion Criteria**

To be eligible for randomisation into this study, each patient must fulfil the following criteria:

1. Male or female,  $\geq 18$  years of age at the time of consent.
2. Admitted to hospital due to the severity of their COVID-19.
3. Positive virus test for SARS-CoV-2 using a validated molecular assay or validated antigen assay.

Patients who had positive virus test for SARS-CoV-2 prior to hospitalisation will be randomised no later than 48 hours after hospital admission. If the virus test was performed more than 96 hours prior to hospitalisation, the test will have to be repeated in the hospital prior to randomisation. Only patients whose repeated virus test is positive will be randomised, no later than 48 hours after confirmation of SARS-CoV-2 infection.

Patients who had positive virus test for SARS-CoV-2 after hospitalisation will be randomised no later than 48 hours after confirmation of SARS-CoV-2 infection.

4. Require oxygen therapy via nasal prongs or mask (OSCI score of 4).
5. Provided informed consent.
6. Female patients must be  $\geq 1$  year post-menopausal, surgically sterile, or using a highly effective method of contraception. Acceptable highly effective methods of contraception include;
  - bilateral tubal occlusion
  - intrauterine device (provided coils are copper-banded)
  - levonorgestrel intrauterine system (e.g., Mirena™)
  - medroxyprogesterone injections (e.g., Depo-Provera™)
  - etonogestrel implants (e.g., Implanon™, Norplan™)
  - normal and low dose combined oral pills
  - norelgestromin/ ethinylestradiol transdermal system
  - intravaginal device (e.g., ethinylestradiol and etonogestrel), desogestrel (e.g., Cerazette™)
  - total sexual abstinence (defined as refraining from heterosexual intercourse)
  - vasectomised sexual partner.

Women should have been stable on their chosen method of birth control for a minimum of 3 months before entering the trial and should continue with birth control for 1 month after the last dose of inhaled IFN-β1a/matching placebo. In addition to the highly effective method of contraception (except for the practice of total sexual abstinence), a condom (in UK with spermicides) should be used by the male partner for sexual

intercourse from randomisation (Visit 2) and for 1 month after the last dose of inhaled IFN- $\beta$ 1a/matching placebo to prevent pregnancy.

7. Women not of childbearing potential are defined as women who are either permanently sterilized (hysterectomy, bilateral oophorectomy, or bilateral salpingectomy), or who are postmenopausal. Women will be considered post-menopausal if they have been amenorrhoeic for 12 months prior to the planned date of randomisation without an alternative medical cause. The following age specific requirements apply:
  - Women <50 years old will be considered post-menopausal if they have been amenorrhoeic for 12 months or more following cessation of exogenous hormonal treatment and if follicle stimulating hormone (FSH) levels are in the postmenopausal range.
  - Women  $\geq$ 50 years old will be considered post-menopausal if they have been amenorrhoeic for 12 months or more following cessation of all exogenous hormonal treatment.

If, in the setting of the pandemic, the use of an acceptable birth control method is not possible, the decision to enrol a woman of childbearing potential should be based on the benefit-risk for the patient, which should be discussed with the patient at the time of the informed consent.

### 3.1.2 Exclusion Criteria

A patient must not be randomised into the study if they meet any of the following criteria:

1. Evidence of ongoing SARS-CoV-2 infection for more than 3 weeks, confirmed by a validated molecular assay or validated antigen assay.
2. Non-invasive ventilation or high-flow oxygen (OSCI score of 5).
3. Mechanical ventilation (continuous or intermittent CPAP or intubation) or admission to intensive care (OSCI score of  $\geq$  6).
4. Previous SARS-CoV-2 infection confirmed by a validated molecular assay or validated antigen assay.
5. Any condition, including findings in the patient's medical history or in the pre-randomisation study assessments that in the opinion of the Investigator, constitute a risk or a contraindication for the participation of the patient into the study or that could interfere with the study objectives, conduct or evaluation.
6. Participation in previous clinical trials of SNG001.
7. Current or previous participation in another clinical trial where the patient has received a dose of an Investigational Medicinal Product (IMP) containing small molecules within 30 days or 5 half-lives (whichever is longer) prior to entry into this study or containing biologicals within 3 months prior to entry into this study.
8. Inability to use a nebuliser with a mouthpiece.
9. Inability to comply with the requirements for storage conditions of study medication in the home setting.
10. History of hypersensitivity to natural or recombinant IFN- $\beta$  or to any of the excipients in the drug preparation.
11. Females who are breast-feeding, lactating, pregnant or intending to become pregnant.
12. Previous SARS-CoV-2 vaccination.

## **4 STUDY MEDICATION**

### **4.1 Allocation to Study Medication**

Patients will be randomised to one of two groups (SNG001 or placebo) in a 1:1 ratio to receive SNG001 (two syringes [Arm 1]), or placebo (two syringes [Arm 2]), according to a pre-specified randomisation schedule in addition to Standard of Care (SOC).

Should a patient withdraw after randomisation, either prior to or after receiving any study medication, the randomisation number will not be re-assigned and the patient will not be replaced.

### **4.2 Breaking the Blind**

The study will be patient- and investigator-blinded with regard to SNG001 or placebo but not the dose. Blinding codes should only be broken in emergency situations for reasons of patient safety. In the event of a need to unblind the patient the investigator (or delegate) will be able to access the patient randomisation list 24 hours a day.

A patient's assignment should only be unblinded in the case of an emergency where the knowledge of the study medication may influence further care of the patient. If a treatment code is unblinded for any reason, the Investigator will notify the Sponsor and a record will be kept of who unblinded the code, the reason for doing so, and the date and time the unblinding occurred. Every attempt will be made to minimise the number of people unblinded. Specific unblinding procedures and instructions can found in the Investigator Site File.

### **4.3 Stopping Rules**

#### **4.3.1 Individual Patient Stopping Rules**

An individual patient will be discontinued permanently from the study medication for any one of the following reasons:

- The patient withdraws consent for study medication only.
- The patient is experiencing intolerable AEs.
- A protocol deviation has occurred that is deemed by the Investigator to put the patient at increased risk.
- The patient is not capable of taking the study medication.
- The Investigator/Sponsor feels that it is in the patient's best interest to be withdrawn from the study medication.
- Pregnancy.
- Other AEs reflecting a safety concern.
- The patient misses more than 4 consecutive doses (i.e. 5<sup>th</sup> dose missed).
- Other (reason to be specified by the Investigator in the electronic case report form [eCRF]).

An individual patient will be discontinued permanently from the study i.e. from study treatment and data collection, for any one of the following reasons:

- The patient withdraws consent for taking part in the study.
- The patient is lost to follow-up.

#### **4.3.2 Trial Stopping Rule**

The trial will be terminated prematurely for any of the following reasons:

- In the opinion of the Sponsor or DSMC/Independent Data Monitoring Committee (IDMC), an unacceptable risk to the safety and welfare of patients is posed by the continuation of the study in light of review of the safety data.
- If clinically significant safety concerns arise within this trial or in any other trial with SNG001 which may impact on the wellbeing of patients in this trial.
- At individual sites, if the study procedures are not being performed according to Good Clinical Practice (GCP) and cannot be remedied or this is a serious breach.
- If recruitment is slow, at the Sponsor's discretion.
- Other (reason to be specified by the Sponsor).

If the trial is terminated prematurely, the Sponsor will ensure that adequate consideration is given to the protection of the patients' interests.

### **4.4 Description of Investigational Medicinal Product and Device**

#### **4.4.1 SNG001**

The drug substance, recombinant IFN- $\beta$ 1a, is manufactured under Good Manufacturing Practice (GMP) conditions by Rentschler Biotechnologie GmbH, Erwin-Rentschler-Straße 21, 88471 Laupheim, Germany. The drug product is manufactured under GMP conditions. The study medication is presented as a ready-to-use aqueous solution (neutral pH).

The IB contains further details of the composition of SNG001.

#### **4.4.2 Placebo**

The placebo will be the same formulation as the study medication but without IFN- $\beta$ 1a (i.e. only the excipients of the SNG001 solution). The IB contains further details of the composition of the placebo.

#### **4.4.3 Inhalation Device**

The study medication will be delivered using the Ultra nebuliser. The Ultra nebuliser is a portable mesh nebuliser marketed by Aerogen. The Aerogen Ultra nebuliser uses palladium vibrating mesh technology, Aerogen Vibronic® is an aerosol drug delivery system. The central aperture plate is 5 mm in diameter and is perforated with 1000 precision formed holes, that vibrate at 128,000 times per second, to produce the optimum particle size for deep lung penetration.

#### **4.5 Dosage and Administration**

While hospitalised, all patients will take their doses of study medication under the supervision of study staff.

SNG001 nebuliser solution is presented in glass syringes containing 0.65 mL of drug product solution containing 12 MIU/mL of IFN- $\beta$ 1a. The Ultra device should be filled with the contents of two syringes (if randomised to Arm 1).

The placebo nebuliser solution is presented in glass syringes containing 0.65 mL of solution containing only the excipients of the SNG001 solution. The Ultra device should be filled with the contents of two syringes (if randomised to Arm 2) (See Appendix A).

Patients will inhale study medication once a day for 14 days.

The study medication should be removed from the fridge for a minimum of 15 minutes prior to taking the dose and for no longer than two hours.

*For patients who have been discharged from hospital during the 14-day treatment phase:*

Patients will have been trained before discharge from hospital to fill the device with IMP and use it. Study staff will call (via telephone or video call) discharged patients on a daily basis and will be able to address any problems and answer any questions patients have about using the device.

If, after multiple training attempts, patients are not capable of taking the study medication at home, even with the support of study staff via phone or video link, they should be withdrawn from the study medication only. Study data should continue to be collected as per the study schedule.

##### **4.5.1 Dose Delivery Times**

It is anticipated that a dose of study medication will take 2 to 6 minutes to be delivered, depending on the patient's inhalation/exhalation breathing pattern.

##### **4.5.2 Packaging and Labelling**

SNG001 and placebo will be presented in glass syringes with needles and plungers as ready-to-use aqueous solutions. Syringes and outer containers will be labelled in accordance with regulatory requirements.

## **4.6 Preparation, Handling and Dispensing**

### **4.6.1 Preparation Time**

The study medication will not require reconstitution and will be supplied in pre-labelled syringes. Study medication will be put into the Ultra nebuliser immediately prior to dosing.

### **4.6.2 Timing of Doses**

The study medication should be taken once daily at approximately the same time of day. There must be a gap of at least 8 hours between doses (see Section 4.9 for the treatment of overdose).

### **4.6.3 Missed or Incomplete Doses**

If a dose is missed or the complete dose is not taken, the following procedure must be followed:

1. Record in the study source document that the dose has been missed/not completed.
2. Continue with any remaining doses as normal (see Section 5.5 for patient withdrawal criteria).

## **4.7 Storage of Study Medication**

Prior to dispensing, the study medication should be stored in a locked and temperature-controlled environment. The study medication should be stored between +2°C and +8°C. A daily temperature log should be kept to ensure the study medication is kept between these temperatures.

Any deviations from the storage conditions should be reported immediately to the Sponsor/Clinical Research Associate (CRA), and the study medication should be placed in quarantine and not used until authorisation has been given by the Sponsor to do so.

Patients who are discharged from hospital during the 14-day treatment phase should be instructed to keep the study medication in a refrigerator. Due to the stability data for the IMP, it is not necessary to keep a record of the fridge storage temperature at the patient's home.

Training will be given to the patients by study staff regarding storage of study medication, disposal of the syringes after they have been used to fill the device, what to do once they have completed all doses and any drug accountability that needs to be completed by the patient.

The study medication can be transported to the patient's home without the need for refrigeration. On arrival at the patient's home, the study medication should be placed in the refrigerator as soon as possible.

## **4.8 Drug Accountability**

The Sponsor will provide the required documentation for study medication accountability, including reconciliation of drugs and maintenance of drug records.

The Investigator must maintain study medication accountability throughout the course of the study in accordance with regulatory requirements.

All patients who take the study medication whilst at home will either be asked to complete a drug accountability form after each dose of study medication, or will be asked questions regarding it by study staff during their daily contact.

The amount of study medication received and dispensed to patients should be documented. Study medication dispensed to a patient should not be returned if unused. Patients will be given instructions for its disposal.

Drug accountability records will include:

- Confirmation and inventory of study medication delivery to the study site.
- An inventory of the study medication that has been provided to each patient.
- The serial number from the controller component of the Ultra nebuliser device.
- Confirmation of administration of each dose of study medication.
- The return to the Sponsor or alternative disposition of unused products.

Dates of use, expiry dates, quantities, batch numbers, and the patient's randomisation and unique patient identification number should also be included on the accountability form.

Unused study medication must not be discarded without authority from the Sponsor or used for any purpose other than the present study.

During the study, where possible due to the nature of the study, the Sponsor (or delegate) will periodically review the drug accountability forms and check these against the actual study medication. Once the study has finished and the drug accountability forms have had a final check, the Sponsor (or delegate) will make arrangements for the return of the study medication, or will authorise their destruction by the study site.

#### **4.9 Treatment of Overdose**

An overdose is defined as any dose greater than the once-a-day dose that is being used in this study i.e. two doses taken less than 8 hours apart. Any overdose must be recorded as a protocol deviation in the study source documents and eCRF.

In the event of overdose, if the patient is not in hospital, hospitalisation should be considered (if applicable) for observation and appropriate supportive treatment should be given. If the patient is in hospital, they should be observed and appropriate supportive treatment should be given. The Sponsor (or delegate) should be contacted in an expedient manner, whether associated with an AE (serious or non-serious) or not, to decide if the patient should continue in the study.

If the patient is experiencing AEs that may relate to an overdose of study medication, the IB should be consulted for details of any specific actions to be taken.

## **5 STUDY PROCEDURES**

### **5.1 Patient Numbering**

Every patient that gives informed consent will be logged and allocated a unique patient identification number. The unique patient identification number will consist of 9 digits (2 for the study number, 2 for the country, 2 for the site number and 3 for patient number), will be assigned sequentially to each patient within each site and will identify the patient throughout the study.

### **5.2 Screening Failures**

Screening failures are defined as patients who have given informed consent for the study, but who have not met the study Inclusion and/or the Exclusion Criteria. Re-screening of patients will be at the discretion of the Medical Monitor. If a patient is re-screened, they will be given a new patient identification number and must give informed consent again.

### **5.3 Study Visits**

#### **5.3.1 Pre-Treatment**

*The following assessments should be performed on Day 0 prior to the first dose preferably in the following order:*

- Informed consent
- Medical history
- COVID-19 infection history
- Patient demographics
- Physical examination (chest exam only unless a physical examination is deemed necessary by the Investigator)
- OSCI
- BCSS
- Vital signs (to include temperature, respiratory rate, heart rate and blood pressure)
- NEWS2
- EQ-5D-5L (complete questionnaire)
- Height and weight
- Urine pregnancy test, if applicable
- Check the patient will be able to use the nebuliser
- Eligibility criteria
- Randomisation
- Blood sample for immunogenicity test (all sites/patients, where collection is feasible)

- Recording of AEs, SAEs and concomitant medications

Eligibility to enter the study may be aided with data captured from SOC procedures and tests performed as part of the admission process.

Patients may take the first dose of study medication either on the same day as the pre-treatment assessments or the next day depending on timings. Day 1 is the day the first dose of study medication is taken. If Day 1 is the same as the day the pre-treatment assessments are performed then the Day 1 assessments do not need to be repeated.

### **5.3.2 Days 1 to 14**

The following assessments should be performed daily from Day 1 to 14.

*For patients that remain in hospital the following assessments should be performed preferably in the following order:*

- Physical examination (chest exam only unless a physical examination is deemed necessary by the Investigator)
- OSCI
- BCSS
- Vital signs (to include daily temperature, respiratory rate, heart rate and blood pressure)
- NEWS2
- EQ-5D-5L (complete questionnaire on Day 7 only. Mobility, self-care and usual activities dimensions from Day 1 to Day 14)
- COVID-19 symptom assessment
- Device training (if required)
- Administration of study medication
- Recording of AEs, SAEs and concomitant medications.

If procedures such as physical exam and vital signs have already been assessed on the day of the visit, this data can be used on the study to avoid the same procedures from being conducted twice.

*For patients who have been discharged from hospital the following should be conducted over telephone/video call preferably in the following order:*

- OSCI
- BCSS
- EQ-5D-5L (complete questionnaire on Day 7 only. Mobility, self-care and usual activities dimensions from Day 1 to Day 14)
- COVID-19 symptom assessment

- Device training (if required)
- Administration of study medication
- Recording of AEs, SAEs and concomitant medications

*If the patient has discontinued dosing\*, attempts should be made to collect the following key data:*

- OSCI
- BCSS
- EQ-5D-5L (complete questionnaire on Day 7 only. Mobility, self-care and usual activities dimensions from Day 1 to Day 14)
- COVID-19 symptom assessment
- Recording of AEs, SAEs and concomitant medications

*\* providing the patient consents to data collection*

### **5.3.3 End of Treatment**

These assessments should be performed on Day 15 (+1 day) (see Sections 4.3.1 and 5.5).

*For patients that remain in hospital the following assessments should be performed, preferably in the following order:*

- Physical examination (chest exam only unless a physical examination is deemed necessary by the Investigator).
- OSCI
- BCSS
- Vital signs
- NEWS2
- EQ-5D-5L (complete questionnaire)
- COVID-19 symptom assessment
- Long-COVID-19 assessment
- Recording of AEs, SAEs and concomitant medications

If procedures such as physical exam and vital signs have already been assessed on the day of the visit, this data can be used on the study to avoid the same procedures from being conducted twice.

*For patients who have been discharged from hospital the following should be conducted over telephone/video call preferably in the following order:*

- OSCI
- BCSS
- EQ-5D-5L (complete questionnaire)
- COVID-19 symptom assessment
- Long-COVID-19 assessment
- Recording of AEs, SAEs and concomitant medications.

#### **5.3.4 Follow-up**

All follow-up assessments can be either done face-to-face if the patient remains in hospital or via telephone/video call if the patient has been discharged from hospital as per the study schedule.

##### **Days 16 to 28**

OSCI, NEWS2, BCSS, AEs, SAEs, the mobility, self-care and usual activities dimensions of EQ-5D-5L and concomitant medications should continue to be collected on a daily basis from the patient.

In addition, complete EQ-5D-5L questionnaire and long-COVID-19 assessments should be performed on Day 28.

NEWS2 will only be assessed if the patient remains in hospital. If the patient has been discharged from hospital NEWS2 will not be assessed. For hospitalised patients, if the components of NEWS2 have already been assessed on the day of the visit, this data can be used on the study to avoid the same procedures being conducted twice.

##### **Days 29 to 35**

OSCI, AEs, SAEs, the mobility, self-care and usual activities dimensions of EQ-5D-5L and concomitant medications should continue to be collected on a daily basis until (and including) Day 35.

##### **Days 36 to 90**

Blood sample for immunogenicity testing should be obtained from the patient (all sites/patients, where collection is feasible) on Day 44 (+/- 5 days).

In addition, BCSS, EQ-5D-5L (complete questionnaire) and long-COVID-19 assessments should be performed on Day 60 and Day 90 (+/- 2 days).

#### **5.4 Unscheduled Visits**

The Investigator can request any of the study procedures described in the study schedule to be repeated as extra follow-up visits or to be added to a study visit where necessary. Unscheduled visits may be necessary if the patient is experiencing prolonged symptoms which the Investigator feels could be related to their study medication or if the Investigator is concerned

about the patient's health. Unscheduled visits/study procedures should be captured in the study source documents.

## **5.5 Patient Withdrawal**

Patients may withdraw from the study at any time at their own request. If a patient does not return for a scheduled visit, every effort should be made to contact the patient. In any circumstance, every effort should be made to document patient outcome, if possible. The Investigator should enquire about the reason for withdrawal, request that the patient disposes of all unused study medication (as instructed) and request the patient undergoes a final visit, if applicable, and follow-up with the patient regarding any unresolved AEs.

Please refer to Section 4.3.1 for individual patient stopping rules.

If patient is withdrawn from the study or from the study medication, the reason for withdrawal will be documented in the study source documents. If the patient is withdrawn from taking the study medication only but continues to give consent for data to be collected, the study schedule should be followed until Day 90.

### **Lost to follow-up:**

Patients will be considered as lost to follow-up if no contact is made with the patient for 5 consecutive visits.

For further guidance, see Appendix B.

## **6 STUDY ASSESSMENTS**

### **6.1 Informed Consent**

Informed consent will be obtained from the patient after the study has been fully explained and the patient has been given ample time to consider the study and discuss it in detail. The method of obtaining consent may vary due to the nature of this study. The method of obtaining and documenting the informed consent and the contents of the consent will comply with International Conference on Harmonisation Guideline for Good Clinical Practice (ICH GCP), and all applicable laws and regulations. Different methods, including written, verbal or electronic consent may be used. Each method will be discussed and approved by the Sponsor (or delegate) before the site takes the first consent. Whichever method is used informed consent will always be obtained from the patient prior to any study related procedures being performed. Informed consent must be obtained by a qualified medical doctor or nurse who have received study training.

If there is a new version of the patient information sheet and informed consent form, patients will be reconsented at their next study visit or, if they were already discharged from the hospital, they may be reconsented verbally over the phone/video link, or electronically. Patients will be informed as soon as possible of the changes and the Sponsor will determine whether the patients need to visit the site prior to their next study visit.

### **6.2 Medical History**

The patient's medical history will be recorded in the study source documents by the Investigator or their delegate to ensure the patient's eligibility.

The medical history should include, but is not limited to;

- Significant current and past medical history in the last 5 years, including hospitalisations, operations and medical history that affects inclusion and exclusion criteria (including sterilisation and contraception).
- Social history regarding smoking habits, alcohol consumption, drug allergies and medication use.
- For women, a menstrual history and history of contraception (if applicable).
- Vaccinations within 30 days prior to randomisation.

### **6.3 COVID-19 Infection History**

The patient's COVID-19 infection history will be recorded in the study source documents by a medical doctor or qualified nurse from the study team.

The study source documents detail questions to be asked and information to be recorded.

### **6.4 Patient Demography**

Patient demographics will be recorded in the study source documents. These will include age, sex, race/ethnicity and residence in care home.

### **6.5 Physical examination**

A physical examination will be performed on Day 1 prior to the first dose by a medical doctor.

Physical examinations may be repeated at various time points throughout the study, as clinically indicated.

If sites have difficulty in completing the chest exam due to risk of contamination/local policies, this will not be considered a protocol deviation, but will need to be explained in a file note prior to the site first patient visit.

Height and weight will be recorded pre-treatment.

The study source documents provide guidance on the physical examination. Results of the examinations will be recorded in the study source documents.

## 6.6 Ordinal Scale for Clinical Improvement

The OSCI to be used in this trial is the 18th February 2020 version as recommended by the WHO.

### Ordinal Scale for Clinical Improvement

Patient State	Descriptor	Score
<i>Uninfected</i>	No clinical or virological evidence of infection	0
<i>Ambulatory</i>	No limitation of activities	1
	Limitation of activities	2
<i>Hospitalised</i>	Hospitalised, no oxygen therapy	3
	Oxygen by mask or nasal prongs	4
<i>Hospitalised</i>	Non-invasive ventilation or high-flow oxygen	5
	Intubation and mechanical ventilation	6
	Ventilation + additional organ support – pressors, RRT, ECMO	7
<i>Dead</i>	Death	8

RRT – renal replacement therapy; ECMO - extracorporeal membrane oxygenation

The assessment should be carried out once a day at approximately the same time each day from Pre-treatment until Day 35. It is recommended that this assessment should be carried out by a clinically qualified member of the study team, i.e. a medical doctor or a qualified nurse. The assessment may be carried out face to face or over the phone/video link.

Patients' status on the day of hospital discharge should be assessed as ambulatory/uninfected (OSCI score of 0, 1 or 2).

After hospital discharge, patients should be asked the following questions about their clinical status and return to the pre-COVID level of activity:

- “In the past 24 hours, did you experience any signs or symptoms of your coronavirus infection?” “Yes” or “No” answer will be required.
- “In the past 24 hours, did you feel that your usual activities (e.g. work, study, housework, family or leisure activities) have returned to the level from before your coronavirus infection and did not require additional assistance/support\*?” “Yes” or “No” answer will be required.

\* assistance/support is defined as additional help of other people and/or requirement for supplemental oxygen (or a higher level of supplemental oxygen), compared to the pre-COVID state.

In order to minimise any potential influence on the patients, the interviewers will be required to read the questions to patients verbatim and not to change or interpret them in any way. The following scoring algorithm will be applied:

<b>Signs/symptoms of coronavirus infection (or virological evidence of infection)?</b>	<b>Limitation of activities?</b>	<b>OSCI Score</b>
No	No	0
Yes	No	1
No	Yes	2
Yes	Yes	2

The applicable OSCI score and the date should be recorded in the study source documents.

## **6.7 Breathlessness, Cough and Sputum Score (BCSS)**

The BCSS is a daily patient-reported outcome measure that was designed as a daily diary in which patients are asked to record the severity of three symptoms: breathlessness, cough and sputum (11).

Each symptom is represented by a single item which is evaluated on a 5-point scale ranging from 0-4, with higher scores indicating more severe symptoms. A mean decline of 1 point on the BCSS total scale signifies a substantial reduction in symptom severity.

This assessment should be carried once a day from Day 1 until Day 28 and then on Day 60 and Day 90 (see Appendix C).

- This assessment should be carried out at approximately the same time each day.
- This assessment should be completed by the patient where possible, however, if needed, the site staff can read out the questions to the patient either face-to-face or over the telephone/video link.
- For patients discharged home during treatment, this assessment will either be recorded on paper by the patient or site staff will ask the patient the questions by telephone/video link.
- This is a patient reported outcome, so for patients that are unable to complete the BCSS due to their medical condition i.e. they are receiving ventilation, data should not be entered into the study worksheet, but the reason for non-completion should be recorded in the study source documents.
- The applicable score (number) for each question should be recorded in the study source documents as well as the date, time and method of assessment.

## **6.8 Vital Signs**

Vital signs include temperature, respiratory rate, heart rate and blood pressure should be recorded daily whilst the patient is hospitalised. When the patient is discharged home vital sign assessments will not be recorded.

Blood pressure and pulse should be measured in the supine position and it is recommended that the patient should have rested quietly for at least 3 minutes prior to the measurement.

## **6.9 National Early Warning Score 2 (NEWS2)**

The data required for NEWS2 will be collected by the site staff and NEWS2 will be calculated by the Sponsor.

NEWS2 is a tool developed by the Royal College of Physicians, which improves the detection and response to clinical deterioration in adult patients and is a key element of patient safety and improving patient outcomes (12).

NEWS2 is based on a simple aggregate scoring system in which a score is allocated to physiological measurements, already recorded in routine practice, when patients present to, or are being monitored in hospital.

Six simple physiological parameters form the basis of the scoring system:

1. Respiration rate
2. Oxygen saturation
3. Any supplementary oxygen
4. Temperature
5. Systolic blood pressure
6. Heart rate
7. Alert, Voice, Pain, Unresponsive.

NEWS2 will be recorded daily whilst the patient is hospitalised. When the patient is discharged home NEWS2 will not be recorded.

## **6.10 Quality of life measured using EQ-5D-5L**

The EQ-5D-5L health-related quality of life questionnaire will be completed on Day 0, Day 7, Day 15, Day 28, Day 60 and Day 90 for all patients (see Appendix D).

In addition, the mobility, self-care and usual activities dimensions of the questionnaire will be assessed daily from Day 1 to Day 35.

The EQ-5D-5L provides a simple descriptive profile and a single index value for health status. The EQ-5D-5L self-rated questionnaire includes a visual analogue scale (VAS), which records the respondent's self-rated health status on a graduated (0–100) scale, with higher scores for

higher health-related quality of life. It also includes the EQ-5D-5L descriptive system, which comprises 5 dimensions of health: mobility, self-care, usual activities, pain/discomfort, and anxiety/depression. The responses record five levels of severity (no problems/slight problems/moderate problems/severe problems/extreme problems) within a particular EQ-5D dimension.

### **6.11 COVID-19 symptom assessment**

The presence of COVID-19 symptoms will be assessed as per the study schedule.

The study source documents detail questions to be asked and information to be recorded.

### **6.12 Assessment of Long-COVID-19 symptoms**

Assessment of long-COVID-19 symptoms will be performed on Day 15, Day 28, Day 60 and Day 90. The following patient reported outcome measures will be used:

- General Anxiety Disorder 7 Questionnaire (GAD-7)
- Patient Health Questionnaire-9 (PHQ-9)
- FACIT Fatigue Scale (Version 4)
- Brief Pain Inventory (Short Form)

The questionnaires are included in Appendix E.

### **6.13 Samples for Laboratory Assessment**

#### **6.13.1 Urine Pregnancy Test**

Early detection pregnancy strip (minimum detection 10 MIU human chorionic gonadotrophin) will be used on Day 0.

Urine pregnancy testing is not required for women who are not of child-bearing potential. Women not of childbearing potential are defined as women who are either permanently sterilized (hysterectomy, bilateral oophorectomy, or bilateral salpingectomy), or who are postmenopausal.

Women will be considered postmenopausal if they have been amenorrhoeic for 12 months prior to the planned date of randomisation without an alternative medical cause. The following age specific requirements apply:

- Women <50 years old would be considered postmenopausal if they have been amenorrhoeic for 12 months or more following cessation of exogenous hormonal treatment and if FSH levels are in the postmenopausal range. If the FSH result is not available at the time of randomisation, the patient must have a negative pregnancy test and agree to use approved contraceptive method. The patient must have been using this contraceptive method for a period of 3 month prior to randomisation.

- Women  $\geq 50$  years old would be considered postmenopausal if they have been amenorrhoeic for 12 months or more following cessation of all exogenous hormonal treatment.

### **6.13.2 Blood sample for immunogenicity testing**

A blood sample will be taken as per the study schedule for immunogenicity testing. This will be completed in all sites and for all patients, where collection of samples is feasible. If a site is not able to participate in immunogenicity sample collection, this will not preclude them from participation in the study. The sample will require processing to obtain serum which is required for the analysis. A laboratory manual describing the requirements for the blood sample and the processing of the sample will be provided separately by the Sponsor (or delegate).

### **6.14 Concomitant Medications**

Medications taken by the patient at the time of informed consent should be recorded on the study worksheet. After this, patients will be asked at each visit if they have taken any new concomitant medications or changed any current medications. Alternatively, information can be obtained from the patient's hospital prescription record. The dosage information, dates of administration and reasons for use will be recorded.

Any other medications which are considered necessary for the patient's welfare should be given at the discretion of the Investigator.

Special care will be taken in questioning the patients on any possible self-medication.

SOC medications for the treatment of COVID-19 (e.g. remdesivir or dexamethasone) can be administered, as required. SOC should be recorded as concomitant medications.

Oxygen therapy will be recorded as concomitant medication.

### **6.15 Prohibited Medications**

Interferons have been reported to reduce the activity of hepatic cytochrome P450 dependent enzymes in humans and animals. Caution should be exercised with patients who are taking medicinal products that have a narrow therapeutic index and are largely dependent on the hepatic cytochrome P450 system for clearance (e.g. some classes of antidepressants). Investigators are able to enrol these patients to the study but should be aware of the possible contraindication.

Patients should be advised to refrain from receiving vaccinations during the treatment and follow up phase up to Day 35, as these could interfere with study assessments.

## **7 ADVERSE EVENT REPORTING**

### **7.1 Adverse Events**

An AE is defined as any untoward medical occurrence in a clinical trial patient during the study; the event does not necessarily have a causal relationship with that treatment. An AE can therefore be any unfavourable and unintended sign (including any clinically significant abnormal laboratory finding), symptom or disease, temporally associated with the use of a medicinal product.

AEs may include the following:

- Apparently unrelated illnesses, including the worsening of a pre-existing illness.
- Injury or accidents.

Patients will be asked about AEs at every study visit. General, non-directed questioning will be used to elicit reports of AEs. If a patient is seen by a physician not involved with the study in relation to an AE, the Investigator should make every effort to contact the treating physician in a timely manner in order to obtain all information necessary to facilitate appropriate reporting of the event.

- The terms ‘COVID-19’ or ‘SARS-CoV-2’ should not be recorded as an AE.
- Individual symptoms related to COVID-19/SARS-CoV-2 infection (e.g. fever, breathlessness, fatigue) present at study entry should not be reported as AEs.
- Worsening of symptoms related to COVID-19/SARS-CoV-2 infection should be reported as AEs if, in the investigators’ opinion, the symptoms are clinically significant or lead to treatment.
- New symptoms related to COVID-19/SARS-CoV-2 infection that start after the initiation of study treatment should be reported as AEs.

### **7.2 Treatment-Emergent Adverse Events**

A treatment-emergent adverse event (TEAE) is an AE that is new in onset, or is a pre-existing condition that is aggravated in severity or frequency, and occurs after the administration of the first dose of study medication. TEAEs include any clinically significant change from baseline readings in any of the study assessments.

Any abnormal result from study assessments will be recorded as an AE, if they result in study withdrawal, are associated with accompanying symptoms, lead to treatment, lead to further diagnostic tests, or is considered by the Investigator to be of clinical significance. These should be recorded in the study source documents.

### **7.3 Serious Adverse Event**

An SAE, according to ICH GCP, is any AE that meets any of the following outcomes:

- Results in death.
- Is life-threatening (i.e. at immediate risk of death).
- Requires in-patient hospitalisation or prolongation of existing hospitalisation.
- Results in persistent or significant disability or incapacity.
- Consists of a congenital anomaly or birth defect.

AEs that may not result in death, may not be life-threatening, or do not require hospitalisation may be considered SAEs when, based upon appropriate medical judgment, they may jeopardise the patient or may require medical or surgical intervention to prevent one of the outcomes listed above. These events must be reported in the same manner as SAEs.

Deterioration of a patient's condition requiring admission to the intensive care unit or mechanical ventilation (whichever happens first) must be reported as an SAE.

Elective and pre-planned surgeries will not be reported as SAEs. Such procedures should be recorded as part of the medical history.

### **7.4 Assessment of Adverse Event**

The Investigator will assess every AE to determine if it is serious or not (see Section 7.3 for definition criteria). This assessment will determine the reporting procedures to be followed.

#### **7.4.1 Relationship to Study Medication**

Investigators (or delegates) need to assess the causality, severity (intensity) and expectedness between the study medication and/or concomitant medication and the AE; this information will be recorded in the study source documents.

##### **7.4.1.1 Causality**

Causality is the relationship between the study medication and the occurrence of the AE. Determination of causality is based on the Investigator's clinical judgment regarding the likelihood that the study medication caused the AE and may include consideration of some or all of the following factors:

- Alternative possible causes of the AE, including the patient's underlying disease or co-morbid conditions, other drugs, other host and environmental factors.
- The temporal sequence between the study medication exposure and the AE.
- Whether the clinical or laboratory manifestations of the AE are consistent with known actions or toxicity of the study medication.

Investigators need to assess the relationship of AEs to the study medication by consulting the IB and using the following definitions:

<b>Definitely related:</b>	A causal relationship of the onset of the event relative to administration of the study medication and there is no other cause to explain the event.
<b>Probably related:</b>	A causal relationship is clinically/biologically reasonable relative to the administration of the study medication and the event is more likely explained by exposure to the study medication than by other factors or causes.
<b>Possibly related:</b>	A causal relationship is clinically/biologically reasonable relative to the administration of the study medication, but the event could have been due to another equally likely cause.
<b>Unlikely to be related:</b>	A causal relationship is considered unlikely to be related to use of the study medication if there are factors (evidence) explaining the occurrence of the event (e.g., progression of the underlying disease, concomitant medication more likely associated with the event) or a convincing alternative explanation for the event.
<b>Unrelated:</b>	A causal relationship is not reasonably related in time to the administration of the study medication, or exposure to the study medication has not occurred.

#### 7.4.1.2 Severity

Severity (intensity) should be assessed according to the following definitions:

<b>Mild:</b>	The patient is aware of the event or symptom, but the event or symptom is easily tolerated, causing minimal discomfort to the patient. The event or symptom should not interfere with everyday activities.
<b>Moderate:</b>	The patient experiences sufficient discomfort to interfere with or reduce normal everyday activities, but responds to symptomatic therapy or rest.
<b>Severe:</b>	Significant impairment of functioning: the patient is unable to carry out normal everyday activities despite symptomatic therapy and/or the patient's life is at risk from the AE.

Note the distinction between the gravity and the severity (intensity) of an AE. Severe is a measure of intensity: thus, a severe reaction is not necessarily a serious reaction. For example, a headache may be severe in intensity, but would not be classified as serious unless it met one of the criteria for SAEs listed above.

## **7.5 Discontinuation of Study Medication due to an AE**

Study medication may be discontinued in response to an AE at any time at the discretion of the patient and/or the Investigator. The Sponsor (or delegate) will be informed within 24 hours of any patient for whom this action is taken.

## **7.6 Pregnancy**

Pregnancy itself is not an AE unless there is a suspicion that the study medication may have interfered with the effectiveness of a contraceptive medication. However, if a female patient becomes pregnant during the conduct of the study, study medication will be discontinued immediately, and the Sponsor (or delegate) should be notified within 24 hours using the Pregnancy Report Form. Follow-up information regarding the outcome of the pregnancy and any foetal or neonatal sequelae should be obtained and documented. For all pregnant patients, on behalf of the Sponsor, study staff will discuss referral for specialist counselling on the possible risks to the unborn baby, and arrangements will be offered to monitor the health of both the patient and the unborn baby.

## **7.7 Adverse Events Recording and Reporting**

### **7.7.1 Reporting of Adverse Events**

All AEs and TEAEs will be reported from the day informed consent is obtained until 28 days after the last administration of the study medication. This includes AEs reported as a result of study procedures.

All AEs and TEAEs occurring during the study assessed by the Investigator (or delegate) as definitely, probably or possibly related to the study medication are to be followed up until resolved or judged to be no longer clinically significant, or until they become chronic or stable to the extent that they can be fully characterized (all follow-up results are to be reported to the sponsor or delegated Clinical Research Organisation [CRO] if requested).

### **7.7.2 Recording of Adverse Events**

Timely, accurate, and complete reporting and analysis of safety information from clinical trials is crucial for the protection of patients, Investigators, and the Sponsor, and is mandated by regulatory agencies. The Sponsor (and delegated CRO) has established standard operating procedures (SOPs) in conformity with regulatory requirements to ensure appropriate reporting of safety information. The process for reporting AEs and SAEs at each site will be assessed by the Sponsor to ensure full compliance with the regulatory requirements.

The Investigator is to report all directly observed AEs and all AEs spontaneously reported by the trial patient using concise medical terminology.

All AEs and TEAEs, regardless of seriousness, severity, or causal relationship to study medication will be recorded in the study source documents. The following items are to be included:

Description of event: whenever possible, signs and symptoms due to a common aetiology will be reported as an integrated diagnosis: e.g. cough, runny nose, sneezing, sore throat, and head congestion would be reported as “upper respiratory infection”. The diagnosis or description will be as specific and complete as possible (i.e. “lower extremity oedema”, rather than just “oedema”).

If a medical condition is known to have caused the injury or accident (e.g. a fall secondary to dizziness), the medical condition (dizziness) and the accident (fall) should be reported as two separate AEs.

In cases of surgical or diagnostic procedures, the condition/illness leading to the procedure is considered as the AE rather than the procedure itself.

In case of a fatality, the cause of death is considered as the SAE, and death is considered as its outcome.

## **7.8 Reporting of Serious Adverse Events**

All AEs assessed as serious by the Investigator (or delegate) according to the criteria specified should be reported and recorded on a SAE form.

SAEs should be reported by the Investigator (or delegated person) to the Sponsor (or delegate), using the SAE report form provided, within 24 hours of a member of the site staff becoming aware of the event, whether or not the event is considered to be related to study medication. This initial report can be made by telephone or in writing. If the initial SAE report is made by telephone, a written report signed by the Investigator (or delegate) must be submitted within 24 hours or the next working day.

All SAEs will be reported from the day informed consent is obtained until 28 days after the last administration of the study medication. SAEs are to be followed until resolution regardless of whether the patient is still participating in the study.

Resolution means:

- The patient has returned to their baseline condition or the Investigator does not expect any further improvement or worsening of the event, or
- The SAE is deemed medically insignificant by the Investigator, or
- The SAE is resolved with residual effects.

**Details of the contact numbers for SAE reporting can be found in the Investigator’s Site File.**

In addition to informing the Sponsor, the Investigator at the study site should notify the host organisation (e.g. Research and Development [R&D]) or any other department as required by local regulations if applicable.

The Sponsor will be responsible for informing the following within the stipulated reporting timelines:

- The Chief Investigator and site investigators
- Institutional Review Board/Independent Ethics Committee (IRB/IEC), and
- The appropriate local regulatory authority

The sponsor will comply with country-specific regulatory requirements relating to safety reporting to the regulatory authority, IRB/IEC, and investigators.

Copies of all correspondence relating to the reporting of SAEs should be maintained in the Investigator Site File and provided to the Sponsor (or delegate). The Investigator will be informed by the Sponsor (or delegate) of any SAEs from other Investigators or clinical studies which have safety implications.

## **8 DATA ANALYSIS/STATISTICAL METHODS**

Data will be entered into a database. The Sponsor (or delegate) will be responsible for data processing and analysis, in accordance with the appropriate data management and analysis procedures.

Analysis methods for key endpoints are described below. Further details on all analyses will be described in the Statistical Analysis Plan (SAP).

### **8.1 Sample Size Determination**

Success will be determined if at least one of the primary endpoints is declared statistically significant by the primary analysis. A sample size of 610 patients in total using a 1:1 randomisation ratio (305 patients per treatment arm) has been chosen to provide at least 90% power to detect a hazard ratio of 1.45 in time to hospital discharge and a hazard ratio of 1.7 in time to recovery and at least 95% power to declare statistical significance on at least one of the primary endpoints. This sample size has been calculated using a global 2-sided alpha level of 0.05 and adjusting for the Hochberg procedure to allow for multiple comparisons. This also allows for an interim analysis to assess futility after the first 300 randomised patients have completed the Day 35 visit or have withdrawn from the study.

The sample size calculation assumes a hospital discharge rate in the placebo treatment arm of 70% at Day 28, a recovery rate in the placebo treatment arm of 30% at Day 28 and a dropout rate of 25% over the 28-day evaluation period with time to dropout exponentially distributed.

By their definition the two primary endpoints are naturally correlated. This correlation has been estimated from prior study data and has also been considered in the sample size calculation.

More details regarding the sample size calculation used for the study will be provided in the SAP.

The key secondary endpoint of progression to severe disease or death within the 35-day study period will be formally tested, provided statistical significance is declared in both of the primary endpoints.

## 8.2 Randomisation

This is a double-blind study. Patients will be randomised to one of two treatment groups (SNG001 or placebo). A pre-specified randomisation schedule will be generated using a 1:1 randomisation ratio.

Arm 1: Approximately 305 patients<sup>^</sup> - SNG001 - contents of 2 syringe<sup>\*</sup> per dose

Arm 2: Approximately 305 patients<sup>^</sup> - Placebo - contents of 2 syringes<sup>+</sup> per dose

<sup>^</sup>Sample size to be confirmed in a protocol amendment.

<sup>\*</sup>Each syringe contains 0.65mL of SNG001 nebuliser solution containing 12 MIU/mL interferon- $\beta$ 1a

<sup>+</sup>Each syringe contains 0.65mL of formulation buffer

## 8.3 Analysis Populations

The intention-to-treat (ITT) population is defined as all randomised patients.

The safety population will include all randomised patients who receive at least one dose of study drug. Patients will be analysed as treated and not per randomised treatment.

The per protocol population will include all patients in the ITT population who do not have any protocol deviations with an impact on efficacy on or prior to Day 35.

## 8.4 Estimands and Intercurrent Events

### 8.4.1 Primary Efficacy Estimands

The first primary estimand will be the hazard ratio of SNG001 versus placebo for the time to hospital discharge of patients in the ITT population. Intercurrent events will be handled as follows:

Intercurrent event	Policy
Death	Composite variable: Patients who die on or prior to Day 28 will have time to hospital discharge right censored at 28 days (the maximum time to discharge).
Premature treatment discontinuation and missed doses	Treatment policy: The time to hospital discharge will be used regardless of treatment discontinuation and the number of missed doses.
Withdrawal from study following hospital discharge	Treatment policy: The time to hospital discharge will be used regardless of study withdrawal and regardless of the duration of follow-up following hospital discharge. This assumes relapse is independent to the treatment effect.
Withdrawal from study prior to hospital discharge	While-on-study: Time to hospital discharge will be right censored at the time of withdrawal from the study or at 28 days, whichever occurs first. The evaluation period for this endpoint is 28 days, therefore estimating the time to

Intercurrent event	Policy
	hospital discharge after Day 28 would not be appropriate. Premature study withdrawal is expected to be independent to the treatment effect, therefore a censoring approach in this scenario is considered appropriate. However, this assumption will be tested by appropriate sensitivity analyses.
Other protocol deviations and concomitant medications affecting efficacy	Treatment policy: Time to hospital discharge will be used regardless of any concomitant medication use or protocol deviations.

A tipping point analysis will be performed to implement a treatment policy strategy for patients who withdraw from the study prior to hospital discharge and prior to Day 28 by imputing missing discharge times. The imputation will be repeated by adjusting the likelihood of hospital discharge in each treatment arm independently, to varying degrees. The first primary analysis will be performed on each of the imputed datasets to test the sensitivity to missing data.

A further sensitivity analysis of the first primary estimand will test the missing at random assumption by imputing missing data after study withdrawal through a missing not at random approach.

A supportive analysis will estimate the hazard ratio of SNG001 versus placebo for the time to hospital discharge of patients in the per protocol population. Intercurrent events will be handled as follows:

Intercurrent event	Policy
Death	Composite variable: Patients who die during the study period will have time to hospital discharge right censored at 28 days (the maximum time to discharge).
Missed 2 of the first 3 scheduled doses	Hypothetical (estimand for a situation with treatment compliance): The time to hospital discharge will be considered missing if two or more doses of the randomised treatment were missed within the first three days of therapy. A single dose within the first three scheduled dosing days is not expected to be sufficient for the study drug to be effective, therefore this approach will help establish the true treatment effect.
Other premature treatment discontinuation	Treatment policy: The time to hospital discharge will be used regardless of treatment discontinuation.
Withdrawal from study following hospital discharge	Treatment policy: The time to hospital discharge will be used regardless of study withdrawal and regardless of the duration of follow-up following hospital discharge. This assumes relapse is independent to the treatment effect.

<b>Intercurrent event</b>	<b>Policy</b>
Withdrawal from study prior to hospital discharge	While-on-study: Time to hospital discharge will be right censored at the time of withdrawal from the study or at 28 days, whichever occurs first. The evaluation period for this endpoint is 28 days, therefore estimating the time to hospital discharge after Day 28 would not be appropriate. Premature study withdrawal is expected to be independent to the treatment effect, therefore a censoring approach in this scenario is considered appropriate.
Other protocol deviations and concomitant medications affecting efficacy	Hypothetical (estimand for a situation with no protocol deviations or concomitant medications affecting efficacy): To establish the true treatment effect the time to hospital discharge will be considered missing if protocol deviations or medications affecting efficacy are recorded on or prior to Day 35.

The second primary estimand will be the hazard ratio of SNG001 versus placebo for the time to recovery of patients in the ITT population. Intercurrent events will be handled as follows:

<b>Intercurrent event</b>	<b>Policy</b>
Death	Composite variable: Patients who die during the study period will have time to recovery right censored at 28 days (the maximum possible time to recovery).
Premature treatment discontinuation and missed doses	Treatment policy: The time to recovery will be used regardless of treatment discontinuation and the number of missed doses.
Withdrawal from study following recovery	Treatment policy: The time to recovery will be used regardless of study withdrawal and regardless of the duration of follow-up following recovery. This assumes relapse is independent to the treatment effect.
Withdrawal from study prior to recovery	While-on-study: Time to recovery will be right censored at the time of withdrawal from the study or at 28 days, whichever occurs first. The evaluation period for this endpoint is 28 days, therefore estimating the time to recovery after Day 28 would not be appropriate. Premature study withdrawal is expected to be independent to the treatment effect, therefore a censoring approach in this scenario is considered appropriate. However, this assumption will be tested by appropriate sensitivity analyses.
Other protocol deviations and concomitant	Treatment policy: Time to recovery will be used regardless of any concomitant medication use or protocol deviations.

Intercurrent event	Policy
medications affecting efficacy	

A tipping point analysis will be performed to implement a treatment policy strategy for patients who withdraw from the study prior to recovery and prior to Day 28 by imputing missing recovery times. The imputation will be repeated by adjusting the likelihood of recovery in each treatment arm independently, to varying degrees. The second primary analysis will be performed on each of the imputed datasets to test the sensitivity to missing data.

A further sensitivity analysis of the second primary estimand will test the missing at random assumption by imputing missing data after study withdrawal through a missing not at random approach.

A supportive analysis will estimate the hazard ratio of SNG001 versus placebo for the time to recovery of patients in the per protocol population. Intercurrent events will be handled as follows:

Intercurrent event	Policy
Death	Treatment policy: Patients who die during the study period will have time to recovery right censored at 28 days (the maximum possible time to recovery).
Missed 2 of the first 3 scheduled doses	Hypothetical (estimand for a situation with treatment compliance): The time to recovery will be considered missing if two or more doses of the randomised treatment were missed within the first three days of therapy. A single dose within the first three scheduled dosing days is not expected to be sufficient for the study drug to be effective, therefore this approach will help establish the true treatment effect.
Other premature treatment discontinuation	Treatment policy: The time to recovery will be used regardless of treatment discontinuation.
Withdrawal from study following recovery	Treatment policy: The time to recovery will be used regardless of study withdrawal and regardless of the duration of follow-up following recovery. This assumes relapse is independent to the treatment effect.
Withdrawal from study prior to recovery	While-on-study: Time to recovery will be right censored at the time of withdrawal from the study or at 28 days, whichever occurs first. The evaluation period for this endpoint is 28 days, therefore estimating the time to recovery after Day 28 would not be appropriate. Premature study withdrawal is expected to be independent to the treatment effect, therefore a censoring approach in this scenario is considered appropriate.

Intercurrent event	Policy
Other protocol deviations and concomitant medications affecting efficacy	Hypothetical (estimand for a situation with no deviations or concomitant medication affecting efficacy): To establish the true treatment effect the time to recovery will be considered missing if protocol deviations or medications affecting efficacy are recorded on or prior to Day 35.

#### 8.4.2 Key Secondary Efficacy Estimands

The first key secondary estimand will be the difference in probabilities for SNG001 versus placebo for progression to severe disease or death within 35 days of first dose of patients in the ITT population. Intercurrent events will be handled as follows:

Intercurrent event	Policy
Death	Composite variable: The definition of severe disease includes patients who die within 35 days of first dose, regardless of their OSCI assessments.
Premature treatment discontinuation and missed doses	Treatment policy: Severe disease will be determined regardless of treatment discontinuation and the number of missed doses.
Withdrawal from study	While-on-study: Patients will only be considered having severe disease if observed and recorded whilst participating in the study. This assumes the treatment effect is independent to reason for study withdrawal.
Other protocol deviations and concomitant medications affecting efficacy	Treatment policy: Severe disease will be determined regardless of any concomitant medication use or protocol deviations.

A tipping point analysis will be performed to implement a treatment policy strategy for patients who withdraw from the study prior to progressing to severe disease and prior to Day 35 by imputing severe disease for patients with missing data. The imputation will be repeated by adjusting the probability of severe disease in each treatment arm independently and recalculating the differences in proportions for each imputed dataset.

A further sensitivity analysis will test the missing at random assumption by imputing severe disease for patients with missing data through a missing not at random approach. Sensitivity analyses will also be conducted only including patients who completed Day 28 and only including patients who withdrew from the study prior to Day 35.

A supportive estimand for the key secondary endpoint will be defined as follows. The difference in probabilities for SNG001 versus placebo for progression to severe disease or death within 35 days of first dose will be estimated for the per protocol population. Intercurrent events will be handled as follows:

Intercurrent event	Policy
Death	Composite variable: The definition of severe disease includes patients who die within 35 days of first dose, regardless of their OSCI assessments.
Missed 2 of the first 3 scheduled doses	Hypothetical (estimand for a situation with treatment compliance): OSCI assessments will be considered missing if two or more doses of the randomised treatment were missed within the first three days of therapy. A single dose within the first three scheduled dosing days is not expected to be sufficient for the study drug to be effective, therefore this approach will help establish the true treatment effect.
Other premature treatment discontinuation	Treatment policy: Severe disease will be determined regardless of treatment discontinuation.
Withdrawal from study	While-on-study: Patients will only be considered having severe disease if observed and recorded whilst participating in the study. This assumes the treatment effect is independent to reason for study withdrawal.
Other protocol deviations and concomitant medications affecting efficacy	Hypothetical (estimand for a situation with no protocol deviations or concomitant medications affecting efficacy): To establish the true treatment effect all OSCI assessments will be considered missing if protocol deviations or medications affecting efficacy are recorded on or prior to Day 35.

The second key secondary estimand will be the difference in probabilities for SNG001 versus placebo for progression to intubation or death within 35 days of first dose of patients in the ITT population. The third key secondary estimand will be the difference in probabilities for SNG001 versus placebo for death within 35 days of first dose of patients in the ITT population. For both the second and third key secondary endpoints the same policies will be used for handling intercurrent events as progression to severe disease or death.

## 8.5 Statistical Analysis Methodology for Efficacy Endpoints

A SAP will be completed prior to study unblinding and database lock.

All collected data will be summarized using descriptive statistics. Summary statistics will consist of the number of observations, mean, standard deviation, minimum, median, and maximum for continuous parameters, and number and percent of subjects for categorical parameters. Changes respective to Baseline will also be evaluated where applicable. Baseline is defined as the last assessment prior to first dose.

### 8.5.1 Analysis of the Primary Efficacy Endpoint

The same method of analysis will be used for both primary endpoints. The primary analysis will use a Cox proportional hazards model. Time to hospital discharge and time to recovery

will be determined using data collected up to Day 28 only. OSCI data collected after Day 28 will only be used to assess relapse for patients who were discharged or recovered prior to Day 28. Relapse will be defined as an OSCI score of  $>2$  for hospital discharge and an OSCI score of  $>1$  for recovery. Patients who die will be administratively censored at Day 28, i.e. time to hospital discharge and time to recovery will be fixed at 28 days for these subjects. Patients who neither recover nor die will be censored at the date of their last OSCI assessment. Covariates will be included for treatment arm, age (as a continuous variable), sex, body mass index (BMI) ( $<30$ ,  $\geq 30$ ), baseline comorbidities, race (White, Non-White), duration of prior symptoms ( $<10$  days,  $\geq 10$  days for hospitalised patients only) and smoking status (Current Smoker, Not Current Smoker). The hazard ratio for recovery of SNG001 versus placebo, 95% confidence intervals (CI) and p-value will be presented.

Ties in event times will be handled using the exact method, or the Efron approximation if computational difficulties arise.

In addition, Kaplan-Meier estimates of time to hospital discharge and time to recovery for 25%, 50% (median time) and 75% of patients will be presented.

### **8.5.2 Analysis of the Key Secondary Efficacy Endpoints**

For the first key secondary endpoint of progression to severe disease or death, a logistic regression model will be applied. The difference in the probability of severe disease progression between SNG001 and placebo will be presented alongside the 95% CIs and p-values. The covariates used for the primary analysis will also be used for the key secondary analysis.

Progression to intubation or death and death will be analysed using the same methods described for the first key secondary endpoint, provided enough events are observed to make the analysis possible.

### **8.5.3 Analysis of Secondary Efficacy Endpoints**

All analyses of secondary endpoints described below will include the covariates used for the primary analysis, unless stated otherwise.

Recovery and hospital discharge will be analysed at Days 7, 14, 21 and 28 using a logistic regression model to determine the difference in proportions for SNG001 versus placebo.

Improvement on the OSCI scale will be analysed at Days 7, 14, 21 and 28 assessed using an ordered logistic regression model assuming proportional odds to estimate the odds ratios for SNG001 versus placebo. Adjacent OSCI categories may be merged for some or all visits if the outcome frequency for some OSCI levels is low for any treatment arm.

Observed OSCI scores will also be summarised descriptively.

Change from baseline in breathlessness and cough scores according to the BCSS and total BCSS score will be assessed using a mixed model for repeated measures. The difference between SNG001 and Placebo across the treatment period (up to Day 14) will be estimated

alongside the 95% CIs and p-values. The covariates used for the primary analysis will be used with an additional covariate for baseline score.

#### **8.5.4 Multiple Testing Strategy**

A Hochberg procedure will be used to test the primary endpoints. Therefore, if the two-sided p-value for the estimate of time to hospital discharge is below 0.05 and the two-sided p-value for the estimate of time to recovery is below 0.05 the null hypotheses for both endpoints will be rejected. Otherwise, if the smallest of the two-sided p-values is below 0.025 only the null hypothesis associated with the smallest p-value will be rejected. In any other circumstance neither null hypothesis will be rejected.

The first key secondary endpoint will be formally tested using a two-sided alpha of 0.05 only if both primary endpoints are statistically significant. The second and third key secondary endpoints will be formally tested, in sequence, using a two-sided alpha of 0.05 only if all prior hypothesis tests are statistically significant.

#### **8.5.5 Handling of Missing Data**

For analyses using the OSCI, including the primary and key secondary endpoints, subjects with an OSCI score of 8, indicating death, will have any subsequent missing OSCI assessments imputed as 8. In addition, if other data sources such as AEs indicate a patient has died all missing OSCI scores on and after the date of death will be imputed as 8.

For relevant analyses, missing OSCI assessments for patients who withdraw from the study alive will be imputed using multiple imputation under a missing at random assumption. Selected analyses will be repeated using a missing not at random approach through an imputation model estimated from patients in the placebo group.

For the mixed models with repeated measures (MMRM) of breathlessness, missing breathlessness scores will be imputed as 4 if the OSCI score at the corresponding visit is  $\geq 5$ . Cough and sputum scores at corresponding visits will be considered missing at random and total BCSS scores will be calculated by summing the imputed symptom scores. Missing breathlessness, cough and sputum scores at all other visits will be considered missing at random and will not explicitly be imputed but will be accounted for by the MMRM analysis. Missing not at random approaches may also be considered for sensitivity analyses.

Further details regarding missing data approaches will be given in the SAP.

### **8.6 Safety Analysis**

Safety analyses will be based primarily on AE information which will be summarised descriptively, including summary of treatment-emergent AEs. AEs will be summarised by Medical Dictionary for Regulatory Activities (MedDRA) preferred term and system organ class. Summaries by severity and drug-relatedness will also be presented.

Vital signs will be summarised by time point for each treatment group and overall.

## 8.7 Other Analyses

Tabulations will be provided for completion/withdrawal status, protocol deviations, study populations, demographic and other baseline characteristics, use of concomitant medications, and exposure of and compliance with study drug.

## 8.8 Subgroup Analyses

The following subgroups will be considered for analysis:

- Age category
- Presence of comorbidities
- Current smoking status
- Sex
- Race
- Prior duration of symptoms
- BMI

The details of analyses to be performed for the subgroups will be defined in the SAP.

## 8.9 Interim Analyses

An unblinded interim analysis is planned to assess futility on the first 300 randomised patients and will be conducted once all of these patients have completed the Day 35 visit or have withdrawn from the study. The primary analyses for the two primary endpoints will be conducted for the interim analysis and provided to an IDMC. The go/no go recommendation made by an IDMC will be based on a non-binding criterion described in the table below.

Criterion	Recommendation
1. the Z statistic from the estimate of treatment effect from the analysis for time to discharge is less than 1.4 <i>and</i> 2. the Z statistic from the estimate of treatment effect from the analysis for time to recovery is less than 1.4	Stop the study for futility
Otherwise	Continue the study so that 610 patients in total are randomised.

In addition, summaries and appropriate analyses of death, intubation or death and severe disease or death will be supplied to the IDMC and should be considered in their recommendation. Full details will be provided in an IDMC charter.

## **8.10 Primary Analysis**

A primary analysis will take place once all randomised patients have completed their Day 35 visit or have withdrawn from the study. The primary analysis will be used to assess the study data against the study objectives. All data summaries documented in the SAP will be produced for the primary analysis and will be reported in a CSR, with the exception of any Day 90 summaries. The study will be fully unblinded from the point of the primary analysis. A follow-up analysis will be conducted once all patients have completed the Day 90 visit or have withdrawn from the study.

## **8.11 Data Safety Monitoring Committee**

The DSMC is an independent multidisciplinary group consisting of suitably qualified individuals who, collectively, have experience in the management of patients with respiratory viral infections and/or other respiratory disease, autoimmune disease or clinical pharmacology and in the conduct and oversight of randomised clinical trials.

The DSMC will be responsible for safeguarding the interests of trial patients by assessing study data, as and when required, to provide recommendations about the progression of the trial.

The DSMC will be advisory to the Chief Investigator, the trial sites and any other Sponsor representative. The Chief Investigator and the Sponsor will be responsible for promptly reviewing the DSMC recommendations, to decide whether to continue or terminate the trial and to determine whether amendments to the protocol or changes in study conduct are required.

The DSMC may recommend that tests or procedures may be repeated to facilitate their decision-making process in order to fulfil their role.

The DSMC will be guided by the DSMC Charter. The Charter will define the primary responsibilities of the DSMC, its membership and the purpose and timing of its meetings. The Charter will also provide the procedures for ensuring confidentiality and proper communication and outline the contents of the reports that will be provided to the DSMC.

# **9 QUALITY CONTROL AND QUALITY ASSURANCE**

## **9.1 Quality Control (Monitoring)**

In order to verify that the study is conducted in accordance with the study protocol, ICH GCP, local regulatory requirements and other applicable guidelines, data will be monitored at regular intervals. This will allow the sponsor to ensure that the data is authentic, accurate and complete. Source data will be verified either on site or remotely. If remote source data verification is required, due to its limitations, critical data such as primary efficacy data and important safety data will be its focus and will comply with local and national regulations. The study Monitoring Plan provides further details on specific monitoring activities.

## **9.2 Quality Assurance**

The Sponsor or its designated representative will assess each study site to verify the qualifications of each Investigator and the site staff and to ensure that the site has all of the required equipment. An Investigators Meeting and/or Site Initiation Meeting will occur where among other things the Investigator will be informed of their responsibilities and procedures for ensuring adequate and correct study documentation.

The Investigator is required to complete the study source documents designed to record all observations and other data pertinent to the study for each patient. Study data for each patient that gives informed consent for the study will be entered into the study source documents by study site personnel.

Instances of missing, discrepant, or uninterpretable data will be queried with the Investigator for resolution. Any changes to study data will be made to the study source documents and documented in an audit trail, which will be maintained within the clinical database.

In compliance with ICH GCP and regulatory requirements, the Sponsor, a third party on behalf of the Sponsor, regulatory agencies, IRBs or IECs may conduct quality assurance audits at any time during or following a study. The Investigator must agree to allow regular monitoring of the study according to ICH GCP requirements and other guidelines and local regulations. The Investigator should also agree to allow auditors direct access to all study-related documents including source documents. They must also agree to allocate their time and the time of their study staff to the auditors in order to discuss findings and issues.

## **10 DATA HANDLING AND RECORD KEEPING**

### **10.1 Study Source Documents and Electronic Case Report Form**

If permitted by local regulations, it is preferable that the study source documents provided by the Sponsor (or delegate) are used. The Investigator and their study team will be responsible for entering data into the study source documents. The Principal Investigator is responsible for ensuring the accuracy of the data entered. The study source documents should be completed during the study visit and once completed, site staff should enter the data into the eCRF.

It is possible that source data may be captured in different locations, for example, patient medical records or directly onto the study source documents.

For screen failures, study source documents will not be monitored. The patient's unique identification number, date of screening, date of informed consent and reason for screen failure should be entered into the eCRF. Please note, no other data need be entered into the eCRF for screen failures.

## **10.2 Record Retention**

Essential documents as defined by ICH GCP are not limited to, but include, all signed protocols and any amendment(s), copies of the completed study source documents, signed informed consent forms/other records of informed consent from all patients who consented, hospital records, diary cards and other source documents, R&D and IEC approvals, copies of previous versions of all study documentation and all related correspondence including approved documents, drug accountability records, study correspondence and a list of the patients' names and addresses.

The Investigator and/or Sponsor must retain copies of the essential documents for a minimum of 15 years following the end of the study.

The Investigator will inform the Sponsor of the storage location of the essential documents and of any changes in the storage location should they occur. The Investigator must contact the Sponsor for approval before disposing of any documentation. The Investigator should take measures to prevent accidental or premature destruction of these documents.

## **11 REGULATORY AND ETHICS**

### **11.1 Regulatory and Ethics Considerations**

The study will be conducted in accordance with the current ICH GCP Guidelines, which are consistent with the ethical principles founded in the Declaration of Helsinki, and in accordance with local applicable laws and regulations.

Procedures to be followed will be those documented in the protocol, study site or Sponsor (or delegate's) SOPs, manuals and guidelines.

### **11.2 Competent Authority Approval**

Before the study is initiated at a site, the Sponsor will obtain approval to conduct the study from the Competent Authority in each of the countries in which this study will be performed (e.g. Medicines and Healthcare products Regulatory Agency (MHRA) in the UK).

### **11.3 Institutional Review Board/Independent Ethics Committee and Hospital Board Requirements**

The Sponsor is responsible for ensuring IRB/IEC approval of the study. Approval of the protocol, informed consent form, advertising and any other information presented to, or seen by, potential patients must be obtained from the appropriate IRB/IEC before the study starts. If a substantial amendment is required during the study, IRB/IEC approval must be obtained prior to their implementation. The Principal Investigators are responsible for ensuring that these actions occur.

If appropriate, as well as approval by the IRB/IEC, prior to starting, the study should obtain approval from the site Hospital Board or other relevant body (e.g. R&D department). Once written approval has been obtained from the country's competent authority (e.g. MHRA), the IRB/IEC, and the site's relevant hospital body, and all other necessary documentation has been

received, the Sponsor (or delegate) will release the study drug and patient recruitment can commence.

Similarly, the Investigator/Chief Investigator must inform the relevant hospital body of all substantial amendments and forward to them evidence of IRB/IEC and competent authority (if appropriate) approval. Amendment approval should be received from the relevant hospital body prior to implementation at site.

#### **11.4 Informed Consent and Screening Data**

Patient informed consent forms will be provided by the Sponsor and will be approved by the IRB/REC. Any changes requested by the IRB/IEC must be approved by the Sponsor (or delegate) prior to the documents being used.

Informed consent will be obtained from each patient prior to any study procedures being performed and data reported.

#### **11.5 Patient Confidentiality**

The Investigator must ensure that the patients' anonymity is maintained. On the study source documents and any other documents submitted to the Sponsor, patients should NOT be identified by their names, but by the assigned unique patient identification number only.

The Investigator (or delegate) should keep a confidential log of the names of all patients and the patient numbers that they have been allocated in order to be able to reveal the identity of the patient should this be required for safety purposes. Documents not for submission to the Sponsor (e.g. signed informed consent forms) should be maintained by the Investigator in strict confidence.

#### **11.6 Investigator Compliance**

The Investigator must be familiar with the study protocol, the IB, the conduct of the study and also their responsibilities as an Investigator according to the ICH GCP guidelines.

No modifications to the protocol will be made without the approval of both the Chief Investigator and the Sponsor. Changes that significantly affect the safety of the patients, the scope of the investigation, or the scientific quality of the study (i.e. efficacy assessments) will require IRB/IEC and regulatory authority (where required) notification before implementation, except where the modification is necessary to eliminate an apparent immediate hazard to human patients.

When circumstances require an immediate departure from procedures set forth in the protocol, the Investigator will contact the Sponsor to discuss the planned course of action. If possible, contact will be made before the implementation of any changes. Any departures from protocol will be fully documented in the source documentation and in the eCRF and/or a protocol deviation log.

## **11.7 Progress Reports and Safety Reports**

Progress and Safety reports should be submitted to the IRB/IEC, competent authority and any necessary local site department as per the country specific requirements. It is the Sponsors responsibility to ensure this happens.

## **12 DEFINITION OF END OF TRIAL**

End of Trial is defined as: Last Patient, Last Visit.

## **13 SPONSOR DISCONTINUATION CRITERIA**

### **13.1 Study Sites**

The Sponsor has the authority to stop activity at any individual site at any time during the study.

### **13.2 Entire Study**

The study will be terminated early if, in the opinion of the Sponsor or Chief Investigator, an unacceptable risk to the safety and welfare of patients is posed by the continuation of the study in light of the data generated. Premature termination of this study may also occur because of a regulatory authority decision, or a change in opinion of the IRB/IEC, or on the advice of the DSMC.

If suspension or termination of the study is necessary, the Sponsor will endeavour to provide advance notification to the sites. The Sponsor should promptly inform each Investigator of its decision and then the Investigator (or delegate) should promptly inform the institution (where applicable) and also the IRB/IEC and any necessary local site department, providing them with a detailed written explanation of the termination or suspension. The Investigator (or delegate) should also promptly inform the trial patients and should assure appropriate therapy and follow up for the patients. In such cases, all study data and unused study medication must be returned to the Sponsor.

## **14 PUBLICATION OF STUDY RESULTS**

The Sponsor commits to publish the results of this study in a peer-reviewed journal of its choice. The results may also be published in a poster or oral presentation at one or more scientific meetings, conferences or symposia.

A clinical study report, written in accordance with the ICH E3 Guideline, will be submitted in accordance with local regulations.

Any and all scientific, commercial, and technical information disclosed by the Sponsor in this protocol, or elsewhere, will be considered the confidential and proprietary property of the Sponsor.

The Investigator shall hold such information in confidence and shall not disclose the information to any third party except to the Investigator's employees and staff who have been

made aware that the information is confidential and who are bound to treat it as such and to whom disclosure is necessary to evaluate that information. The Investigator shall not use such information for any purpose other than determining mutual interest in performing the study and, if the parties decide to proceed with the study, for the purpose of conducting the study.

The Investigator understands that the information developed from this clinical study will be used by the Sponsor in connection with the development of SNG001 and other drugs and diagnostics, and, therefore, may be disclosed as required to other Clinical Investigators, business partners and associates and government agencies. The Investigator also understands that, to allow for the use of the information derived from the clinical study, the Investigator has the obligation to provide the Sponsor with complete test results and all data developed in the study.

No publication or disclosure of study results will be permitted except under the terms and conditions of a separate written agreement between the Sponsor and the Investigator and/or the Investigator's institution.

## 15 APPENDICES

### Appendix A: Aerogen Ultra Device

#### Introduction

The Aerogen® Solo System is an iteration of the Aerogen® Pro System. It is intended for hospital use only to nebulise physician-prescribed medications for inhalation which are approved for use with a general purpose nebuliser.

The Aerogen® Ultra is an accessory specific to the Aerogen Solo nebuliser it facilitates intermittent and continuous nebulisation.

The life of the nebuliser has been validated for intermittent use for a maximum of 28 days based upon a typical usage profile of 4 treatments per day.

An appropriate filter is attached to mitigate the level of device related emissions into the atmosphere. The filter should be attached via a connector.

#### Preparing the Ultra for patient use

**Note:** Inspect for device integrity and correct valve placement prior to use.



Insert Aerogen Solo nebuliser firmly into Aerogen Ultra in orientation shown.



### Loading the study medication

1. The study medication should always be kept in the fridge.
2. Remove the box of study medication from the fridge.
3. Remove two syringes of study medication from the box. The syringes should be removed from the fridge for at least 15 minutes prior to taking the dose of study medication and used as soon as possible after this time. The remaining study medication should be returned to the fridge.

**Caution:** The study medication should not be out of the fridge for more than two hours.

4. Open the plug on the Aerogen Solo.
5. Take the first syringe and remove the needle cap. Place the needle into the Aerogen Solo and then being careful not to spill any study medication, push the plunger down, placing all the study medication into the Aerogen Solo. Repeat this process with the second syringe.
6. Ensure that the empty syringes are placed into the sharps bin provided, taking care at all times with the needle.
7. Close the Aerogen Solo plug.
8. Connect the USB controller to the Aerogen Solo port.
9. Connect the Aerogen USB controller to the provided AC/DC Adapter

- **Administering the study medication**

**Note:** If supplemental oxygen is required, nasal cannula can be used during study medication administration. If nasal cannula is required, the Physician or nurse observing the

administration of the dose should ensure that the mist in the chamber is emptying with each breath. This will ensure that the dose is being administered to the patient.

1. The study medication should be taken sitting comfortably in an upright position.
2. Apply the nose clip and ensure a tight seal between the lips and the mouthpiece. Note: a nose clip is not necessary if nasal cannula are required by the patient.
3. To start the nebuliser, press and release the blue ON/OFF power button. The green indicator light illuminates. Aerosol should now be visible, which indicates correct operation.
4. To stop the nebulizer at any time, press the ON/OFF power button. The indicator turns off to indicate that nebulization has stopped.
5. Breathe normally in and out through the mouth. Do not breathe through the nose.
6. Ensure that the cloud of aerosol disappears from the chamber.



**Note:** For optimum use of the Aerogen Solo, ensure it is correctly orientated as shown in the image.

- **Pausing the treatment**

If required, the treatment may be paused by pressing and releasing the blue ON/Off power button on the USB controller for 1 second. Restart the treatment as soon as possible.


- **Cleaning the Aerogen Ultra**

To ensure optimal performance of the Aerogen Ultra, nebulise a few drops of normal saline, shake off excess and allow to air dry.

- **WARNINGS AND CAUTIONS**

- Do not cover Aerogen Ultra valves during use.
- Do not autoclave any component of the kit.
- Ensure tubing is safely orientated to prevent strangulation hazard.
- Visually check Aerogen Ultra post-rinsing to ensure that valves have not become dislodged.
- There will be extra tubing (oxygen) in the package.
- Do not use with a closed face mask or a standard oxygen mask.
- Do not exceed recommended oxygen flow for system.

## Appendix B: Withdrawal Flow chart



## **Appendix C: BCSS questionnaire**

### **BCSS questions and responses:**

1. How much difficulty did you have breathing today?
  - 0 = None – unaware of any difficulty
  - 1 = Mild – noticeable when performing strenuous activity (e.g. running)
  - 2 = Moderate – noticeable even when performing light activity (e.g. bedmaking or carrying groceries)
  - 3 = Marked – noticeable when washing or dressing
  - 4 = Severe – almost constant, present even when resting
  
2. How was your cough today?
  - 0 = No cough – unaware of coughing
  - 1 = Rare – cough now and then
  - 2 = Occasional – less than hourly
  - 3 = Frequent – one or more times an hour
  - 4 = Almost constant – never free of cough or need to cough
  
3. How much trouble did you have due to sputum today?
  - 0 = None – unaware of any trouble
  - 1 = Mild – rarely caused trouble
  - 2 = Moderate – noticeable trouble
  - 3 = Marked – caused a great deal of trouble
  - 4 = Severe – almost constant trouble

## **Appendix D: EQ-5D-5L**

Under each heading, please tick the ONE box that best describes your health TODAY

### **MOBILITY**

- I have no problems in walking about ☐
- I have slight problems in walking about ☐
- I have moderate problems in walking about ☐
- I have severe problems in walking about ☐
- I am unable to walk about ☐

### **SELF-CARE**

- I have no problems washing or dressing myself ☐
- I have slight problems washing or dressing myself ☐
- I have moderate problems washing or dressing myself ☐
- I have severe problems washing or dressing myself ☐
- I am unable to wash or dress myself ☐

### **USUAL ACTIVITIES** (*e.g. work, study, housework, family or leisure activities*)

- I have no problems doing my usual activities ☐
- I have slight problems doing my usual activities ☐
- I have moderate problems doing my usual activities ☐
- I have severe problems doing my usual activities ☐
- I am unable to do my usual activities ☐


### **PAIN / DISCOMFORT**

- I have no pain or discomfort ☐
- I have slight pain or discomfort ☐
- I have moderate pain or discomfort ☐
- I have severe pain or discomfort ☐
- I have extreme pain or discomfort ☐

### **ANXIETY / DEPRESSION**

- I am not anxious or depressed ☐
- I am slightly anxious or depressed ☐
- I am moderately anxious or depressed ☐
- I am severely anxious or depressed ☐
- I am extremely anxious or depressed ☐

The best health  
you can imagine



- We would like to know how good or bad your health is TODAY.
- This scale is numbered from 0 to 100.
- 100 means the best health you can imagine.  
0 means the worst health you can imagine.
- Mark an X on the scale to indicate how your health is TODAY.
- Now, please write the number you marked on the scale in the box below.

YOUR HEALTH TODAY:

The worst health  
you can imagine

## Appendix E: Long-COVID-19 patient reported outcome measures

<b>General Anxiety Disorder 7 Questionnaire (GAD-7)</b>				
<b>Over the <u>last two weeks</u>, how often have you been bothered by the following problems?</b>	Not at all	Several Days	More than half the days	Nearly every day
Feeling nervous, anxious or on edge	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Not being able to stop or control worrying	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Worrying too much about different things	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Trouble relaxing	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Being so restless that it's hard to sit still	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Becoming easily annoyed or irritable	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Feeling afraid as if something awful might happen	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Column totals:	<input type="checkbox"/> + ( <input type="checkbox"/> x 2 ) + ( <input type="checkbox"/> x 3 )			
	= Total Score <input type="checkbox"/> <input type="checkbox"/>			
<p>If you checked off any problems, how difficult have these problems made it for you to do your work, take care of things at home, or get along with other people?</p> <p>           Not difficult at all      Somewhat difficult      Very difficult      Extremely difficult  <input type="checkbox"/>                              <input type="checkbox"/>                              <input type="checkbox"/>                              <input type="checkbox"/> </p>				
<p>From the Primary Care Evaluation of Mental Disorders Patient Health Questionnaire (PRIME-MD PHQ). The PHQ was developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke and colleagues. For research information, contact Dr. Spitzer at rls8@columbia.edu. PRIME-MD® is a trademark of Pfizer Inc. Copyright© 1999 Pfizer Inc. All rights reserved.</p>				

<b>Patient Health Questionnaire-9 (PHQ-9)</b>				
<b>Over the <u>last two weeks</u>, how often have you been bothered by any of the following problems?</b>	Not at all	Several Days	More than half the days	Nearly every day
Little interest or pleasure in doing things?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Feeling down, depressed, or hopeless?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Trouble falling asleep or staying asleep, or sleeping too much?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Feeling tired or having little energy?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Poor appetite or overeating?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Feeling bad about yourself – or that you are a failure of have let yourself or your family down?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Trouble concentrating on things, such as reading the newspaper or watching television?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Moving or speaking so slowly that other people could have noticed? Or so fidgety or restless that you have been moving a lot more than usual?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Thoughts that you would be better off dead, or thoughts of hurting yourself in some way?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3
Column totals:	<input type="checkbox"/> + ( <input type="checkbox"/> x 2 ) + ( <input type="checkbox"/> x 3 )			
	= Total Score <input type="checkbox"/> <input type="checkbox"/>			
From the Primary Care Evaluation of Mental Disorders Patient Health Questionnaire (PRIME-MD PHQ). The PHQ was developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke and colleagues. For research information, contact Dr. Spitzer at rls8@columbia.edu. PRIME-MD® is a trademark of Pfizer Inc. Copyright© 1999 Pfizer Inc. All rights reserved.				


<b>FACIT Fatigue Scale (Version 4)</b>					
<b>Below is a list of statements that other people with your illness have said are important. Please circle or mark one number per line to indicate your response as it applies to the past 7 days.</b>					
	<b>Not at all</b>	<b>A little bit</b>	<b>Some -what</b>	<b>Quite a bit</b>	<b>Very much</b>
I feel fatigued	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
I feel weak all over	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
I feel listless (“washed out”)	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
I feel tired	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
I have trouble <u>starting</u> things because I am tired	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
I have trouble <u>finishing</u> things because I am tired	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
I have energy	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
I am able to do my usual activities	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
I need to sleep during the day	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
I am too tired to eat	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
I need help doing my usual activities	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
I am frustrated by being too tired to do the things I want to do	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
I have to limit my social activity because I am tired	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
Copyright 1987, 1997 FACIT.org					

# **Brief Pain Inventory (Short Form)**

1. Throughout our lives, most of us have had pain from time to time (such as minor headaches, sprains, and toothaches). Have you had pain other than these everyday kinds of pain today?

Yes ☐ No ☐

2. On the diagram, shade in the areas where you feel pain. Put an X on the area that hurts the most.



3. Please rate your pain by marking the box beside the number that best describes your pain at its **worst** in the last 24 hours.

☐ 0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10  
No Pain Pain As Bad As You Can Imagine

4. Please rate your pain by marking the box beside the number that best describes your pain at its **least** in the last 24 hours.

☐ 0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10  
No Pain Pain As Bad As You Can Imagine

5. Please rate your pain by marking the box beside the number that best describes your pain on the **average**.

☐ 0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10  
No Pain Pain As Bad As You Can Imagine

6. Please rate your pain by marking the box beside the number that tells how much pain you have **right now**.

☐ 0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10  
No Pain Pain As Bad As You Can Imagine

7. What treatments or medications are you receiving for your pain?										
8. In the last 24 hours, how much relief have pain treatments or medications provided? Please mark the box below the percentage that most shows how much <b>relief</b> you have received?										
0%	10%	20%	30%	40%	50%	60%	70%	80%	90%	100%
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
No Relief									Complete Relief	
9. Mark the box beside the number that describes how, during the past 24 hours, pain has interfered with your:										
A. General Activity										
<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6	<input type="checkbox"/> 7	<input type="checkbox"/> 8	<input type="checkbox"/> 9	<input type="checkbox"/> 10
Does Not Interfere									Completely Interferes	
B. Mood										
<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6	<input type="checkbox"/> 7	<input type="checkbox"/> 8	<input type="checkbox"/> 9	<input type="checkbox"/> 10
Does Not Interfere									Completely Interferes	
C. Walking ability										
<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6	<input type="checkbox"/> 7	<input type="checkbox"/> 8	<input type="checkbox"/> 9	<input type="checkbox"/> 10
Does Not Interfere									Completely Interferes	
D. Normal Work (includes both work outside the home and housework)										
<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6	<input type="checkbox"/> 7	<input type="checkbox"/> 8	<input type="checkbox"/> 9	<input type="checkbox"/> 10
Does Not Interfere									Completely Interferes	
E. Relations with other people										
<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6	<input type="checkbox"/> 7	<input type="checkbox"/> 8	<input type="checkbox"/> 9	<input type="checkbox"/> 10
Does Not Interfere									Completely Interferes	
F. Sleep										
<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6	<input type="checkbox"/> 7	<input type="checkbox"/> 8	<input type="checkbox"/> 9	<input type="checkbox"/> 10
Does Not Interfere									Completely Interferes	
G. Enjoyment of life										
<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6	<input type="checkbox"/> 7	<input type="checkbox"/> 8	<input type="checkbox"/> 9	<input type="checkbox"/> 10
Does Not Interfere									Completely Interferes	
Copyright 1991 Charles S. Cleeland, PhD Pain Research Group All rights reserved										

## 16 REFERENCES

1. Mantlo E, Bukreyeva N, Maruyama J, Paessler S, Huang C. Antiviral activities of type I interferons to SARS-CoV-2 infection. *Antiviral Res.* 2020;179:104811.
2. Agrawal A. Mechanisms and implications of age-associated impaired innate interferon secretion by dendritic cells: a mini-review. *Gerontology.* 2013;59(5):421-6.
3. Singanayagam A, Loo SL, Calderazzo M, Finney LJ, Trujillo Torralbo MB, Bakhsoliani E, et al. Antiviral immunity is impaired in COPD patients with frequent exacerbations. *Am J Physiol Lung Cell Mol Physiol.* 2019;317(6):L893-L903.
4. Wark PA, Johnston SL, Bucchieri F, Powell R, Puddicombe S, Laza-Stanca V, et al. Asthmatic bronchial epithelial cells have a deficient innate immune response to infection with rhinovirus. *J Exp Med.* 2005;201(6):937-47.
5. Epidemiology Working Group for Ncip Epidemic Response Chinese Center for Disease Control Prevention. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. *Zhonghua Liu Xing Bing Xue Za Zhi.* 2020;41(2):145-51.
6. Synairgen announces positive results from trial of SNG001 in hospitalised COVID-19 patients [press release]. 20th July 2020 2020.
7. World Health Organisation. WHO R&D Blueprint novel Coronavirus COVID-19 Therapeutic Trial Synopsis. 2020.
8. Dmitrienko A, Tamhane AC, Wiens BL. General multistage gatekeeping procedures. *Biom J.* 2008;50(5):667-77.
9. Dugernier J, Hesse M, Vanbever R, Depoortere V, Roeseler J, Michotte JB, et al. SPECT-CT Comparison of Lung Deposition using a System combining a Vibrating-mesh Nebulizer with a Valved Holding Chamber and a Conventional Jet Nebulizer: a Randomized Cross-over Study. *Pharmaceutical research.* 2017;34(2):290-300.
10. Tepper JS, Kuehl PJ, Cracknell S, Nikula KJ, Pei L, Blanchard JD. Symposium Summary: "Breathe In, Breathe Out, Its Easy: What You Need to Know About Developing Inhaled Drugs". *International journal of toxicology.* 2016;35(4):376-92.
11. Leidy NK, Rennard SI, Schmier J, Jones MK, Goldman M. The breathlessness, cough, and sputum scale: the development of empirically based guidelines for interpretation. *Chest.* 2003;124(6):2182-91.
12. Royal College of Physicians. National Early Warning Score (NEWS) 2: Standardising the assessment of acute-illness severity in the NHS. Updated report of a working party. . 2017.